

SERINE PROTEASE INHIBITORS

This invention relates to compounds which are inhibitors of serine proteases, to pharmaceutical compositions thereof and to their use in the treatment of the human or animal body.

The serine proteases are a group of proteolytic enzymes which have a common catalytic mechanism characterized by a particularly reactive Ser residue. Examples of serine proteases include trypsin, tryptase, chymotrypsin, elastase, thrombin, plasmin, kallikrein, Complement C1, acrosomal protease, lysosomal protease, cocoonase, α -lytic protease, protease A, protease B, serine carboxypeptidase II, subtilisin, urokinase, Factor VIIa, Factor IXa, and Factor Xa.

The serine proteases have been investigated extensively over a period of several decades and the therapeutic value of inhibitors of serine proteases is well understood.

Serine protease inhibitors play a central role in the regulation of a wide variety of physiological process including coagulation, fibrinolysis, fertilization, development, malignancy, neuromuscular patterning and inflammation. It is well known that these compounds inhibit a variety of circulating proteases as well as proteases that are activated or released in tissue. It is also becoming clear that serine protease inhibitors inhibit critical cellular processes, such as adhesion, migration, free radical production and apoptosis. In addition, animal experiments indicate that intravenously administered serine protease inhibitors, variants or cells expressing serine protease inhibitors, provide a protective effect against tissue damage.

Serine protease inhibitors have also been predicted to have potential beneficial uses in the treatment of disease in a wide variety of clinical areas such as oncology, neurology, haematology, pulmonary medicine, immunology, inflammation and infectious disease.

In particular serine protease inhibitors may be beneficial in the treatment of thrombotic diseases, asthma, emphysema, cirrhosis, arthritis, carcinoma, melanoma, restenosis, atheroma, trauma, shock and reperfusion injury.

5 Thus for example an inhibitor of Factor Xa has value as a therapeutic agent as an anticoagulant, e.g. in the treatment and prevention of thrombotic disorders. The use of a Factor Xa inhibitor as an anticoagulant is desirable in view of the selectivity of its effect. Many clinically approved
10 anticoagulants have been associated with adverse events owing to the non-specific nature of their effects on the coagulation cascade.

Also, there are well-known associations of α_1 protease inhibitor deficiency with emphysema and cirrhosis and C1
15 esterase inhibitor deficiency with angioedema.

It has now been found that certain aromatic compounds are particularly effective as inhibitors of serine proteases, especially proteases with negatively charged P1 specificity pockets, and most especially the serine protease Factor Xa.

20 The Factor Xa inhibitors of this invention are potentially useful for the prophylaxis or treatment of thrombotic disorders such as amongst others venous thrombosis, pulmonary embolism, arterial thrombosis, myocardial ischaemia, myocardial infarction, and cerebral thrombosis. They
25 potentially have benefit in the treatment of acute vessel closure associated with thrombolytic therapy and restenosis, e.g. after transluminal coronary angioplasty or bypass grafting of the coronary or peripheral arteries and in the maintenance of vascular access patency in long term
30 hemodialysis patients.

Factor Xa inhibitors of this invention may, with benefit, form part of a combination therapy with an anticoagulant with a different mode of action or with a thrombolytic agent.


It has been reported in WO99/11658 and WO99/11657 that certain benzamidine and aminoisoquinoline derivatives carrying a bulky lipophilic side chain are excellent inhibitors of serine proteases. Unfortunately, it has since been found that benzamidine compounds of WO 99/11658 in general demonstrate poor oral bioavailability.

Surprisingly, it has now been found that certain other aromatic compounds also show inhibitory activity against serine proteases, in particular Factor Xa, despite the lack of the amidino or 1-aminoisoquinoline functionality previously believed to be crucial for activity as a factor Xa inhibitor. Many of these compounds also possess other structural features that further distinguish them from the compounds of WO99/11658 and WO99/11657.

Where compounds of the invention have been tested, they have generally demonstrated superior oral bioavailability in comparison with benzamidines disclosed in WO 99/11658. Also, it has been found that the compounds of the invention perform excellently in the prothrombin time assay (PT) when compared to aminoisoquinolines of similar factor Xa activity and structure. The PT assay is a coagulation assay and it is widely accepted that direct acting Factor Xa inhibitors which perform well in the PT assay are more likely to be good antithrombotics. Compounds of the invention have also been found to exhibit a good duration of action following oral administration.

In WO99/09053 certain 2-aminobenzamide compounds are disclosed as potential motilin receptor antagonists and in US 3268513 similar 2-aminobenzamide compounds are suggested as potential antibacterial agents. However, the novel compounds of the present invention have not before been suggested as potential serine protease inhibitors.

Thus viewed from an one aspect the invention provides a serine protease inhibitor compound of formula (I)



wherein:

R₂ is a 5 or 6 membered aromatic carbon ring optionally

5 interrupted by a nitrogen, oxygen or sulphur ring atom,
optionally being substituted in the 3 and/or 4 position (in
relation to the point of attachment of X-X) by halo, nitro,
thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano,
haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or
0 difluoromethoxy, carboxy, acyloxy, MeSO_2^- or R_1 , or the
substituents at the 3 and 4 positions taken together form a
fused ring which is a 5 or 6 membered carbocyclic or
heterocyclic ring optionally substituted by halo, haloalkoxy,
haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl,
5 alkynyl or R_{1j} , and optionally substituted in the position
alpha to the X-X group (i.e. 6 position for a six membered
aromatic ring etc) by amino, hydroxy, halo, alkyl, carboxy,
alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio
with the proviso that R_2 cannot be aminoisquinolyl;

20 each X independently is a C, N, O or S atom or a CO,
CR_{1a}, C(R_{1a})₂ or NR_{1a} group, at least one X being C, CO, CR_{1a}
or C(R_{1a})₂;

each R_{1a} independently represents hydrogen or hydroxyl, alkoxy, alkyl, aminoalkyl, hydroxyalkyl alkoxyalkyl, 25 alkoxycarbonyl, alkylaminocarbonyl, alkoxycarbonylamino, acyloxymethoxycarbonyl or alkylamino optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl;

R₁ is as defined for R_{1a}, provided that R₁ is not unsubstituted aminoalkyl;

30 Y (the α -atom) is a nitrogen atom or a CR_{1b} group;

Cy is a saturated or unsaturated, mono or poly cyclic, homo or heterocyclic group, preferably containing 5 to 10 ring

atoms and optionally substituted by groups R_{3a} or phenyl optionally substituted by R_{3a} or $R_{3i}X_i$;


each R_{3a} independently is R_{1c} , amino, halo, cyano, nitro, thiol, alkylthio, alkylsulphonyl, alkylsulphenyl, triazolyl, imidazolyl, tetrazolyl, hydrazido, alkylimidazolyl, thiazolyl, alkylthiazolyl, alkylloxazolyl, oxazolyl, alkylsulphonamido, alkylaminosulphonyl, aminosulphonyl, haloalkoxy, haloalkyl, a group of the formula $-C(X^3)N(R^{11})R^{12}$ (wherein X^3 is O or S; and R^{11} and R^{12} are independently selected from hydrogen, methyl or ethyl or together with the nitrogen atom to which they are attached form a pyrrolidin-1-yl, piperidin-1-yl or morpholino group), or $-OCH_2O-$ which is bonded to two adjacent ring atoms in Cy;

X_i is a bond, O, NH or CH_2 ;

R_{3i} is phenyl pyridyl or pyrimidinyl optionally substituted by R_{3a} ;

R_{1b} , R_{1c} and R_{1j} are as defined for R_{1a} ; and

$-L-Lp(D)_n$ is



q is 1 or 2;

Q is methylene; and R_q is NR_aR_b in which each of R_a and R_b independently is hydrogen or C_{1-3} alkyl; or one of R_a and R_b is hydrogen or methyl and the other of R_a and R_b is (3-6C)cycloalkyl, pyrid-4-yl, $-CH_2-R_c$ or $-CH_2-R_d$ in which R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, $CONH_2$, SO_2NH_2 , methylaminosulphonyl, dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and in which R_d is isopropyl or cyclopentyl, or NR_aR_b is azetidino, pyrrolidino, piperidino, morpholino, thiomorpholino, piperazino, or tetrahydro-1,4-diazepino [in which a pyrrolidino or piperidino may be a

3,4-didehydro derivative and in which a azetidino, pyrrolidino, piperidino, morpholino, thiomorpholino, piperazino, or tetrahydro-1,4-diazepino may be optionally substituted on a ring carbon atom by hydroxy, amino, (1-3C)alkoxy, (1-3C)hydroxyalkyl, (1-3C)alkyl, carboxy, methoxycarbonyl or ethoxycarbonyl (provided that the amino, hydroxy or alkoxy substituent is not on a ring carbon atom which is included in a double bond, or adjacent to a ring oxygen, sulfur or nitrogen atom) and in which the piperazino or tetrahydro-1,4-diazepino may bear a methyl group at the 4-position]; or a physiologically-tolerable salt thereof, e.g. a halide, phosphate or sulphate salt or a salt with ammonium or an organic amine such as ethylamine or meglumine.

In the compounds of the invention, where the alpha atom is carbon it preferably has the conformation that would result from construction from a D- α -aminoacid $\text{NH}_2\text{-CR}_{1b}(\text{Cy})\text{-COOH}$ where the NH_2 represents part of X-X. Likewise the fourth substituent R_{1b} at an alpha carbon is preferably a methyl or hydroxymethyl group or hydrogen. It will be appreciated that the compounds of formula (I) may exist in racemic or chiral form, and that the preferred D-isomer may be administered in a racemic mixture with the L-isomer, or alone.

In the compounds of the invention, unless otherwise indicated, aryl groups preferably contain 5 to 10 ring atoms optionally including 1, 2 or 3 heteroatoms selected from O, N and S; alkyl, alkenyl or alkynyl groups or alkylene moieties preferably contain up to 6 carbons, e.g. C_{1-6} or C_{1-3} ; cyclic groups preferably have ring sizes of 3 to 8 atoms; and fused multicyclic groups preferably contain 8 to 16 ring atoms.

Examples of particular values for R_{1a} are: hydrogen, methyl or ethyl. R_{1a} is preferably a hydrogen atom.

The linker group from the R_2 group to the alpha atom is preferably selected from -CH=CH- , -CONH- , $\text{-CONR}_{1a}\text{-}$, -NH-CO- ,


-NH-CH₂-, -CH₂-NH-, -CH₂O-, -OCH₂-, -COO-, -OC=O- and -CH₂CH₂-. Preferably, the X moiety nearest to the alpha atom is an NH or O atom, most preferably a NH group. The X moiety alpha to the aromatic ring is preferably a carbon based group such as CH₂ or CO, preferably CO. Thus a particularly preferred linker X-X is -CONH-. In an alternative embodiment the linker is preferably a -OCH₂- group.

Examples of particular values for R_{1b} are: hydrogen, (1-4C)alkyl, such as methyl or hydroxy(1-4C)alkyl, such as hydroxymethyl. R_{1b} is preferably a hydrogen atom.

The alpha atom (Y) is preferably a CH or C(CH₃) group, especially CH.

In the group -L-Lp(D)_n, preferably the azetidino, pyrrolidino, piperidino, morpholino, thiomorpholino, piperazino or tetrahydro-1,4-diazepino in NR_aR_b is optionally substituted on a ring carbon atom by methyl, hydroxy or hydroxymethyl.

A preferred sub-group of compounds of formula I is that in which wherein -L-Lp(D)_n is of the formula:



wherein:


q is 1 or 2;

Q is methylene; and R_q is NR_aR_b in which each of R_a and R_b independently is hydrogen or C₁₋₃alkyl; or one of R_a and R_b is hydrogen or methyl and the other of R_a and R_b is -CH₂-R_c or -CH₂-R_d in which R_c is pyridyl or phenyl (which phenyl may bear a fluoro, chloro, methyl, CONH₂, SO₂NH₂, methylaminosulphonyl, dimethylaminosulphonyl, methylsulphonylamino, methoxy or methylsulphonyl substituent) and in which R_d is isopropyl or cyclopentyl, or NR_aR_b is pyrrolidino, piperidino, morpholino, piperazino, or

tetrahydro-1,4-diazepino in which a pyrrolidino or piperidino may be a 3,4-didehydro derivative and in which a pyrrolidino, piperidino, piperazino, or tetrahydro-1,4-diazepino may bear a methyl group at the 4-position.

5 q is preferably 2.

Preferably $-L-Lp(D)_n$ is of the formula:



wherein:

q is 1 or 2;

10 Q is methylene; and R_q is NR_aR_b in which each of R_a and R_b independently is hydrogen or C_{1-3} alkyl; or one of R_a and R_b is hydrogen and the other is (3-6C)cycloalkyl or pyrid-4-yl; or NR_aR_b is azetidino, pyrrolidino, piperidino or piperazino [in which a pyrrolidino or piperidino may be a 3,4-didehydro
15 derivative and in which a azetidino, pyrrolidino, piperidino or piperazino may be optionally substituted on a ring carbon atom by methyl, hydroxy or hydroxymethyl (provided that the hydroxy substituent is not on a ring carbon atom which is included in a double bond, or adjacent to a ring nitrogen
20 atom) and in which the piperazino may bear a methyl group at the 4-position].

More preferably R_q is selected from dimethylamino, diethylamino, prop-2-ylamino, pyrrolidino, 3-pyrrolino, 3-hydroxypyrrolidino, 3-hydroxymethylpyrrolidino, piperidino, 3-
25 hydroxypiperidino, 4-hydroxypiperidino, 4-hydroxymethylpiperidino, piperazino and 4-methylpiperazino.


Examples of particular values for R_q are dimethylamino, diethylamino, prop-2-ylamino, pyrrolidino, 3-pyrrolino, 3-hydroxypyrrolidino, 3-hydroxymethylpyrrolidino, 3(S)-
30 hydroxypyrrolidino, 3(S)-hydroxymethylpyrrolidino, 3(R)-

hydroxymethylpyrrolidino, piperidino, 4-hydroxypiperidino, 4-hydroxymethylpiperidino, 3-hydroxypiperidino, piperazino and 4-methylpiperazino.

When R_a or R_b represents a C_{1-3} alkyl group, this may be, for example, a methyl or ethyl group. When R_b represents a C_{3-6} cycloalkyl group, this may be, for example, a cyclohexyl group.

Preferably R_Q is NR_aR_b in which R_a is hydrogen or C_{1-3} alkyl and R_b is C_{1-3} alkyl; or R_a is hydrogen and R_b is (3-
 10 6C)cycloalkyl or pyrid-4-yl; or NR_aR_b is azetidino, pyrrolidino, piperidino, morpholino, thiomorpholino or piperazino [in which a pyrrolidino, piperidino or piperazino may be optionally substituted on a ring carbon atom by hydroxy or hydroxymethyl (provided that the hydroxy substituent is not
 15 on a ring carbon atom which is adjacent to a ring nitrogen atom) and in which the piperazino may bear a methyl group at the 4-position].

Most preferably, the group $L-Lp(D)_n$ is selected from




wherein:

m represents 0 or 1;

and when R_3 is present as a substituent on a saturated ring, it is selected from hydrogen, hydroxy, amino, (1-3C)alkoxy, (1-3C)hydroxyalkyl, (1-3C)alkyl, carboxy, methoxycarbonyl and ethoxycarbonyl.

5 For example specific groups of formula $L-Lp(D)_n$ include



Cy is preferably an optionally R_{3a} substituted: phenyl, pyridyl, thienyl, thiazolyl, naphthyl, piperidinyl, furanyl, 10 pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, oxazolyl, imidazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, pyrimidinyl, pyridazinyl, quinoloyl, isoquinolyl, benzofuryl, benzothienyl or cycloalkyl group, or a phenyl group substituted by $R_{31}X_1$ in which X_1 is a bond, O, NH or CH_2 and 15 R_{31} is phenyl, pyridyl or pyrimidinyl optionally substituted by R_{3a} .

The cyclic group attached to the alpha carbon may thus be an optionally R_{3a} substituted phenyl, pyridyl (such as pyrid-2-yl, pyrid-3-yl or pyrid-4-yl), thienyl (such as thien-2-yl 20 or thien-3-yl), thiazolyl (such as thiazol-2-yl, thiazol-4-yl or thiazol-5-yl), naphthyl (such as naphth-1-yl), piperidinyl (such as piperidin-4-yl) or cycloalkyl, such as a cyclohexyl group.

Examples of values for Cy when it represents phenyl substituted by $R_{31}X_1$ are 3-(2-pyridyl)phenyl, 3-(3-pyridylphenyl and 3-(4-pyridylphenyl).

Examples of particular values for R_{3a} are:-

hydrogen;
hydroxyl;
30 for alkoxy: methoxy or ethoxy;

- for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or ethyl, or alkylaminoalkyl, such as methylaminomethyl or dimethylaminomethyl;
- 5 for hydroxyalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: hydroxymethyl or carboxy;
for alkoxyalkyl: methoxymethyl;
for alkoxycarbonyl: methoxycarbonyl or ethoxycarbonyl;
- 10 for alkylaminocarbonyl: methylaminocarbonyl or dimethylaminocarbonyl;
for aminoalkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: aminomethyl, CONH_2 or CH_2CONH_2 ;
for alkylamino optionally substituted by hydroxy, alkylamino,
- 15 alkoxy, oxo, aryl or cycloalkyl: (1-6C)alkanoylamino, such as acetylamino;
for alkoxycarbonylamino: methoxycarbonylamino, ethoxycarbonylamino or t-butoxycarbonylamino;
amino;
- 20 for halo: fluoro, chloro or bromo;
cyano;
nitro;
thiol;
for alkylthio: methylthio;
- 25 for alkylsulphonyl: methylsulphonyl or ethylsulphonyl;
for alkylsulphenyl: methylsulphenyl;
for alkylsulphonamido: methylsulphonylamido or ethylsulphonylamido;
for alkylaminosulphonyl: methylaminosulphonyl or
- 30 ethylaminosulphonyl;
aminosulphonyl;
for haloalkoxy: trifluoromethoxy;
for haloalkyl: trifluoromethyl;

for a group of formula $-C(X^3)N(R^{11})R^{12}$: pyrrolidin-1-ylcarbonyl, piperidin-1-ylcarbonyl or morpholin-1-ylcarbonyl; and $-OCH_2O-$ which is bonded to two adjacent ring atoms in Cy.

Examples of particular values for R_{1C} are:

- 5 hydrogen;
hydroxyl;
for alkoxy: methoxy or ethoxy;
for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or
- 10 ethyl, or alkylaminoalkyl, such as methylaminomethyl or dimethylaminomethyl;
for hydroxyalkyl: hydroxymethyl;
for alkoxyalkyl: methoxymethyl;
for alkoxycarbonyl: methoxycarbonyl or ethoxycarbonyl;
- 15 for alkylaminocarbonyl: methylaminocarbonyl or dimethylaminocarbonyl;
for alkoxycarbonylamino: methoxycarbonylamino, ethoxycarbonylamino or t-butoxycarbonylamino;
for alkylamino optionally substituted by hydroxy, alkylamino,
- 20 alkoxy, oxo, aryl or cycloalkyl: (1-6C)alkanoylamino, such as acetylamino; and
for aminoalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: aminomethyl, $CONH_2$ or CH_2CONH_2 .

Preferably R_{3a} is hydrogen, hydroxyl, methoxy, methyl,
25 amino, fluoro, chloro, ethylsulphonylamino, amido or methylaminocarbonyl.

Examples of values for R_{3i} are phenyl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl pyrimid-2-yl and pyrimid-6-yl.

- Examples of particular values for Cy are phenyl, 4-
- 30 aminophenyl, 4-amidophenyl, 4-(N-methyl)amidophenyl, 4-(N,N-dimethyl)amidophenyl, 2-chlorophenyl, 2-methylphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 4-hydroxyphenyl, 2-methoxyphenyl, 4-methoxyphenyl, 4-carboxyphenyl, 3-ethylsulphonylaminophenyl, thien-2-yl, thien-3-yl, thiazol-4-

yl, thiazol-5-yl, 2-methylthiazol-4-yl, pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, piperidin-4-yl, 1-methylpiperidin-4-yl, cyclohexyl and naphth-1-yl. Other examples are:

4-carbamoylphenyl; furan-2-yl; furan-3-yl; imidazol-2-yl;







5 thiazol-2-yl; 2-aminothiazol-4-yl; isoquinolin-5-yl;

isoquinolin-8-yl; quinolin-5-yl; and quinolin-8-yl.

Further examples are: 2-trifluoromethylphenyl; 2-methylthiophenyl; 2-methylsulphonylphenyl; 3-bromophenyl;

3-cyanophenyl; and benzo[b]thiophen-3-yl.

10 Particular mention is made of the following values for Cy:



or



wherein:

X' is selected from O, S and NMe;

X'' is selected from O and S;

5 X''' is selected from O, S, NH and NMe;

Y' is selected from hydrogen, amino and methyl;

R₀ is selected from hydrogen, methyl, fluoro, chloro, trifluoromethyl, methoxy, methylthio, methylsulphinyl and methylsulphonyl;

10 R_m is selected from hydrogen, methyl, fluoro, chloro, trifluoromethyl, methoxy, methylthio, methylsulphinyl, methylsulphonyl, carboxy, methoxycarbonyl and a group of the formula -C(X³)N(R¹¹)R¹² (wherein X³ is O or S, and R¹¹ and R¹² are independently selected from hydrogen, methyl or ethyl or

15 together with the nitrogen atom to which they are attached form a pyrrolidin-1-yl, piperidin-1-yl or morpholino group);

R_p is selected from hydrogen and fluoro; or

R₀ and R_m or R_m and R_p form an -OCH₂O- group; or

R₀ and R_m together with the ring to which they are attached
20 form a 5 or 6 membered aryl or heteroaryl ring (wherein the heteroaryl ring contains 1 or 2 heteroatoms selected from nitrogen, oxygen and sulfur); and

one of R₀₁ and R₀₂ is hydrogen and the other is R₀.

Preferably Cy is selected from phenyl (optionally

25 substituted by ethyl, prop-2-yl, phenoxy, hydroxy, ethoxy, benzyloxy, prop-2-yloxy, nitro, amino, acetylamino, methylsulfonylamino, dimethylamino, chloro, methoxy, trifluoromethyl, methylthio, methylsulfonyl, tert-butylthio, tert-butylsulfonyl, aminosulfonyl or carbamoyl), pyridyl,
30 thienyl, furanyl, imidazolyl, thiazolyl (optionally substituted by amino or methyl), naphthyl, isoquinolinyl and quinolinyl.

Examples of values for Cy when it represents thiazolyl substituted by amino or methyl are 2-aminothiazol-4-yl and 2-

methylthiazol-4-yl.


Preferably Cy is selected from phenyl, 2-chlorophenyl, 2-methoxyphenyl, 4-carbamoylphenyl, pyrid-2-yl, pyrid-4-yl, thien-2-yl, thien-3-yl, furan-2-yl, furan-3-yl, imidazol-2-yl, 5 thiazol-2-yl, thiazol-4-yl, 2-amino-thiazol-4-yl, thiazol-5-yl, naph-1-thyl, isoquinolin-5-yl, isoquinolin-8-yl, quinolin-4-yl, quinolin-5-yl and quinolin-8-yl.

More preferably Cy is selected from phenyl, 2-methoxyphenyl, 4-carbamoylphenyl, pyrid-2-yl, thien-2-yl, 10 thien-3-yl, furan-2-yl, furan-3-yl, imidazol-2-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl and quinolin-4-yl.

A value for Cy of particular interest is phenyl.

Referring to the group R₂, examples of a 5 or 6 membered aromatic carbon ring optionally interrupted by a nitrogen, 15 oxygen or sulphur ring atom are phenyl; pyrrolyl, such as 2-pyrrolyl; pyridyl, such as 3-pyridyl; pyrazinyl, such as 2-pyrazinyl; furyl, such as 2-furyl; and thienyl, such as 2-thienyl or 3-thienyl. Preferably the ring is interrupted (i.e. a carbon atom is replaced) by at most one heteroatom. 20 More preferably the ring is phenyl, 2-thienyl or 2-pyrrolyl. Most preferably, the ring is phenyl.

When the ring is phenyl, the group R₂ may be a group of formula



25 in which R₅ is amino, hydroxy or hydrogen, and R₆ and R₇ which may be the same or different represent halo, nitro, thiol, cyano, haloalkyl, haloalkoxy, amido, hydrazido, amino, alkylthio, alkenyl, alkynyl or R₁ or taken together form a 5

or 6 membered fused carbocyclic ring or 5 membered heterocyclic ring, which may itself be substituted by R_{1j} , amino, halo, cyano, nitro, thiol, alkylthio, haloalkyl, haloalkoxy.

- 5 When the substituents at the 3 and 4 positions taken together form a fused ring which is a 5 or 6 membered carbocyclic or heterocyclic ring, examples of the resultant bicyclic ring are naphthyl, such as 2-naphthyl; benzimidazolyl, such as benzimidazol-5-yl or benzimidazol-6-yl; isoquinolinyl, such as isoquinolin-7-yl; indolyl, such as indol-2-yl, indol-5-yl or indol-6-yl; indazolyl, such as indazol-5-yl; indazol-6-yl; 3,4-methylenedioxyphenyl; dihydroindolyl, such as 2,3-dihydroindol-6-yl; benzothiazolyl, such as benzothiazol-2-yl or benzothiazol-6-yl;
- 10 benzo[b]thiophenyl, such as benzo[b]thiophen-2-yl; benzofuryl, such as benzofur-2-yl; imidazo[1,2-a]pyrimidinyl, such as imidazo[1,2-a]pyrimidin-2-yl; tetrahydroimidazo[1,2-a]pyrimidinyl, such as tetrahydroimidazo[1,2-a]pyrimidin-2-yl; and benzisoxazolyl, such as benzisoxazol-5-yl.
- 15 Preferably R_2 is phenyl, thien-2-yl, naphthyl, indol-2-yl, indol-6-yl, benzo[b]furan-5-yl, benzo[b]thiophen-2-yl or benzimidazol-2-yl, optionally substituted as defined hereinabove.

R_2 preferably represents:

- 25 (i) phenyl optionally being substituted in the 3 and/or 4 position by halo, nitro, thiol, haloalkoxy, hydrazido, alkylhydrazido, amino, cyano, haloalkyl, alkylthio, alkenyl, alkynyl, acylamino, tri or difluoromethoxy, carboxy, acyloxy, MeSO_2 - or R_1 , and optionally substituted at the 6 position by
- 30 amino, hydroxy, halo, alkyl, carboxy, alkoxycarbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio;
- (ii) naphth-2-yl optionally substituted at the 6 or 7 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} and optionally

substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio;

(iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or benzisoxazol-5-yl

5 optionally substituted at the 3 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(iv) benzimidazol-5-yl or benzothiazol-6-yl optionally substituted at the 2 position by amino;

10 (v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;

15 (vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;

(viii) pyrazol-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

20 (ix) pyrid-2-yl optionally substituted at the 5 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(x) pyrid-3-yl optionally substituted at the 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, 25 hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(xi) benzofur-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, 30 hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by alkyl and optionally substituted at the 5 or 6 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j};

(xiii) indol-6-yl substituted at the 5 position by amino, hydroxy, halo (such as fluoro or chloro), alkyl, carboxy, alkoxy, carbonyl, cyano, amido, aminoalkyl, alkoxy or alkylthio and optionally substituted at the 3 position by halo (such as 5 chloro), haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} ; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the 3 position by amino, hydroxy, halo, alkyl, carboxy, cyano, amido, aminoalkyl, alkoxy or alkylthio and at the 5 or 6 10 position by halo, haloalkoxy, haloalkyl, cyano, nitro, amino, hydrazido, alkylthio, alkenyl, alkynyl or R_{1j} .

Examples of particular values for substituents that may be present on R_2 are:

for halo: fluoro, chloro, bromo or iodo;

15 nitro;

thiol;

for haloalkoxy: difluoromethoxy or trifluoromethoxy;

hydrazido;

for alkylhydrazido: methylhydrazido;

20 amino;

cyano;

for haloalkyl: trifluoromethyl;

for alkylthio: methylthio;

for alkenyl: vinyl;

25 for alkynyl: ethynyl;

for acylamino: acetyl amino;

carboxy;

for acyloxy: acetoxy;

hydroxy;

30 for alkyl: methyl or ethyl;

amido (CONH_2);

for aminoalkyl: aminomethyl; and

for alkoxy: methoxy or ethoxy.

Preferably R_2 is optionally substituted by 1 or 2 substituents selected from fluoro, chloro, amino, methyl, ethyl and methoxy.

Examples of particular values for R_1 are:

- 5 hydrogen;
hydroxy;
for alkoxy: methoxy or ethoxy;
for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or
10 ethyl, alkylaminoalkyl, such as dimethylaminomethyl, or alkanoyl, such as acetyl;
for hydroxyalkyl: hydroxymethyl;
for alkoxyalkyl: methoxymethyl;
for alkoxycarbonyl: methoxycarbonyl;
15 for alkylaminocarbonyl: methylaminocarbonyl;
for alkylamino: methylamino, ethylamino or dimethylamino;
for hydroxyalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: carboxyl or carboxymethyl; and
for aminoalkyl substituted by hydroxy, alkylamino, alkoxy,
20 oxo, aryl or cycloalkyl: amido (CONH_2) or amidomethyl.

Examples of particular values for R_{1j} are:

- hydrogen;
hydroxy;
for alkoxy: methoxy or ethoxy;
25 for alkyl optionally substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: alkyl, such as methyl or ethyl, or alkanoyl, such as acetyl;
for hydroxyalkyl: hydroxymethyl;
for alkoxyalkyl: methoxymethyl;
30 for alkoxycarbonyl: methoxycarbonyl;
for alkylamino: methylamino, ethylamino or dimethylamino;
for hydroxyalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: carboxyl or carboxymethyl; and

for aminoalkyl substituted by hydroxy, alkylamino, alkoxy, oxo, aryl or cycloalkyl: amido (CONH_2) or amidomethyl.

More preferably R_2 represents:

- (i) phenyl optionally being substituted in the 3 and/or 5 4 position by fluoro, chloro, bromo, iodo, nitro, difluoromethoxy, trifluoromethoxy, amino, cyano, trifluoromethyl, methylthio, vinyl, carboxy, acetoxy, MeSO_2 -, hydroxy, methoxy, ethoxy, methyl, methoxycarbonyl, methylamino, ethylamino or amido, and optionally substituted 10 at the 6 position by amino, hydroxy, fluoro, methoxycarbonyl, cyano or aminomethyl (preferably phenyl substituted in the 4 position by chloro, amino, vinyl, methylamino, methyl or methoxy, optionally at the 3 position with amino or hydroxy, and optionally at the 6 position with amino or hydroxy);
- 15 (ii) naphth-2-yl optionally substituted at the 6, position by hydroxy and optionally substituted at the 3 position by amino or hydroxy;
- (iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, indazol-5-yl, indazol-6-yl, benzothiazol-6-yl or benzisoxazol-5-yl 20 optionally substituted at the 3 position by chloro, bromo, amino, methyl or methoxy (preferably indol-6-yl optionally substituted at the 3 position by chloro, bromo, methyl or methoxy);
- (iv) benzimidazol-5-yl or benzothiazol-6-yl optionally 25 substituted at the 2 position by amino;
- (v) thien-2-yl or thien-3-yl optionally substituted at the 4 or 5 position by methylthio, methyl or acetyl;
- (vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;
- 30 (vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;
- (viii) pyrazol-2-yl substituted at the 5 position by methyl;

(ix) pyrid-2-yl optionally substituted at the 6 position by chloro;

(x) pyrid-3-yl optionally substituted at the 4 position by chloro;

5 (xi) benzofur-2-yl optionally substituted at the 3 position by chloro, methyl or methoxy, at the 5 or 6 position by methyl and at the 6 position by methoxy;

(xii) indol-2-yl optionally substituted on the indole nitrogen atom by methyl and optionally substituted at the 5 or
10 6 position by fluoro, chloro, bromo, methyl or methoxy;

(xiii) indol-6-yl substituted at the 5 position by chloro, fluoro or hydroxy and optionally substituted at the 3 position by chloro or methyl; or

(xiv) benzo[b]thiophen-2-yl optionally substituted at the
15 3 position by fluoro, chloro or methyl, and optionally substituted at the 5 or 6 position by fluoro, chloro, methyl, hydroxy, or methoxy.









Examples of particular values for R₂ are:

(i) phenyl, 2-aminophenyl, 3-aminophenyl, 2-amino-3-
20 fluorophenyl, 2-amino-4-fluorophenyl, 2-amino-4-chlorophenyl, 2-amino-3-bromophenyl, 2-amino-3-nitrophenyl, 2-amino-4-nitrophenyl, 3,4-dimethoxy-5-aminophenyl, 2-amino-4-methylphenyl, 2-amino-3-methylphenyl, 2-amino-3-methoxyphenyl, 3,4-diaminophenyl, 3,5-diaminophenyl, 3-amino-4-fluorophenyl,
25 3-amino-4-chlorophenyl, 3-amino-4-bromophenyl, 3-amino-4-hydroxyphenyl, 3-amino-4-carboxymethylphenyl, 3-amino-4-methylphenyl, 3-amino-4-methoxyphenyl, 2-fluorophenyl, 4-fluoro-3-cyanophenyl, 3-chlorophenyl, 3-chloro-4-hydroxyphenyl, 3-chloro-5-hydroxyphenyl, 4-chlorophenyl, 4-chloro-2-
30 hydroxyphenyl, 4-chloro-3-hydroxyphenyl, 4-chloro-3-methylphenyl, 4-chloro-3-methoxyphenyl, 4-bromophenyl, 4-bromo-3-methylphenyl, 4-iodophenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 3-cyano-5-aminophenyl, 2-hydroxyphenyl, 2-hydroxy-4-methoxyphenyl, 3-hydroxyphenyl, 3-

- hydroxy-4-methylphenyl, 2,4-dihydroxyphenyl, 3,4-dihydroxyphenyl, 3-hydroxy-4-methoxyphenyl, 4-difluoromethoxyphenyl, 4-trifluoromethoxyphenyl, 4-trifluoromethylphenyl, 4-methylthiophenyl, 4-methoxycarbonylphenyl, 4-acetoxyphenyl, 4-methanesulfonylphenyl, 3-methylphenyl, 3-methyl-5-aminophenyl, 4-methylphenyl, 4-vinylphenyl, 4-methoxyphenyl, 4-ethoxyphenyl, 4-methoxy-3-chlorophenyl, 4-methoxy-3-methylphenyl, 3-methylaminophenyl, 4-methylaminophenyl, 4-ethylaminophenyl or 2-aminomethylphenyl;
- (ii) naphth-2-yl, 3-aminonaphth-2-yl, 3-hydroxynaphth-2-yl or 6-hydroxynaphth-2-yl;
- (iii) isoquinolin-7-yl, indol-5-yl, indol-6-yl, 3-chloroindol-6-yl, 3-bromoindol-6-yl, 3-methylindol-6-yl, 3-methoxyindol-6-yl, indazol-5-yl, 3-aminoindazol-5-yl, indazol-6-yl, benzothiazol-6-yl, 3-aminobenzisoxazol-5-yl;
- (iv) benzimidazol-5-yl, 2-aminobenzimidazol-5-yl, or benzothiazol-6-yl;
- (v) thien-2-yl, 5-methylthien-2-yl, 5-methylthio-thien-2-yl, 5-acetylthien-2-yl or thien-3-yl;
- (vi) 3,4-methylenedioxyphenyl, 2,3-dihydroindol-6-yl, 3,3-dichloro-2-oxo-indol-6-yl or 1-methyl-3-aminoindazol-5-yl;
- (vii) benzothiazol-2-yl, imidazo[1,2-a]pyrimidin-2-yl or tetrahydroimidazo[1,2-a]pyrimidin-2-yl;
- (viii) 5-methylpyrazol-2-yl;
- (ix) 5-chloropyrid-2-yl;
- (x) pyrid-3-yl, 6-chloropyrid-3-yl;
- (xi) benzofur-2-yl, 5-chlorobenzofur-2-yl, 3-methylbenzofur-2-yl, 5-methylbenzofur-2-yl, 6-methoxybenzofur-2-yl;
- (xii) indol-2-yl, 5-fluoroindol-2-yl, 5-chloroindol-2-yl, 5-methylindol-2-yl, 5-methoxyindol-2-yl, 6-methoxyindol-2-yl and 1-methyl-indol-2-yl;
- (xiii) 5-fluoroindol-6-yl; or

(xiv) benzo[b]thiophen-2-yl, 5-chloro- benzo[b]thiophen-2-yl or 6-chlorobenzo[b]thiophen-2-yl.

R_2 may, for example, be selected from one of the formula (A') to (H'):



5

wherein X_4 is O or S, R_{13} is selected from hydrogen, chloro or methyl and R_{14} is selected from hydrogen, methyl, ethyl, fluoro, chloro, and methoxy and R_{15} is selected from hydrogen, methyl, fluoro, chloro and amino.

Preferably R_2 is of the formula (A') (wherein R_{14} is selected from hydrogen, methyl, ethyl, fluoro, chloro, and methoxy and R_{15} is selected from hydrogen, methyl, fluoro, chloro and amino) or of the formula (B') (wherein R_{13} is chloro) or of the formula (C') (wherein R_{13} is selected from

15

hydrogen, methyl and chloro) or of the formula (D') (wherein R_{13} is selected from hydrogen and chloro) or of the formula (E') (wherein R_{13} is hydrogen) or of the formula (G') (wherein R_{13} is chloro).

- 5 More preferably R_2 is 4-methoxyphenyl, 5-chloroindol-2-yl, 3-chloroindol-6-yl, indol-6-yl or 3-methylindol-6-yl.

R_2 is preferably of the formula (A') and R_{14} and R_{15} are as defined hereinabove. More preferably R_2 is of the formula (A') and R_{14} is methoxy and R_{15} is hydrogen.

- 10 It is preferred that at least one of R_6 and R_7 be other than hydrogen and that R_6 , if present, is preferably a substituent containing one or more polar hydrogens such as hydroxy, amino, alkylamino, alkylaminoalkyl, aminocarbonyl, alkylaminocarbonyl, hydrazo and alkylhydrazo; alternatively R_6
15 and R_7 are joined together in the formation of a naphthyl or indolyl or azaindolyl or diazaindolyl group.

- It is especially preferred that R_6 be amino and R_7 be chloro, bromo, methyl, methoxy or vinyl; or that R_6 and R_7 taken together form an indolyl ring with the NH at the 6-
20 position or taken together form a naphthyl ring.


The compounds of the invention may be prepared by conventional chemical synthetic routes or by routes as illustrated by the following examples.

- The compounds of the formula (I) may be prepared by
25 forming the -X-X- bond from appropriate intermediates. For example, when -X-X- is -CONH- or -CO-NR_{1a}-, by reacting a compound of the formula (10): $H_2N-Y-(Cy)-L-Lp(D)_n$ with a compound of the formula R_2-COOH , under conditions known for the formation of an amide bond. The reaction is conveniently
30 carried out in the presence of a benzotriazole-based reagent such as 1-hydroxybenzotriazole or 1-hydroxy-7-azabenzotriazole, in an inert organic solvent such as dimethylformamide and/or methylene chloride. The reaction mixture is usually taken to 0°C and then a dehydrating agent

such as dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide added. Other suitable reagents and solvents are known in the art. For example, an acid of formula R_2COOH may be converted into an acid halide, such as an acid chloride, and then reacted with the compound of formula (10) in the presence of a base, such as pyridine. Another reagent is diethyl cyanophosphonate

Compounds wherein -X-X- is -NHCO- or -NHCH₂- may be formed from the appropriate intermediates using reaction conditions for the formation of an amide bond as described above and if necessary subsequent reduction of the resulting amide bond.

Compounds of the formula (I) wherein -X-X- is of the formula -CH₂NH- may be prepared by reducing the corresponding compound of the formula (I) wherein -X-X- is -CONH-, or by reaction of a compound of formula (10)



with a compound of the formula R_2CHO and reducing the intermediate of formula (I) where X-X is C=N- with, for example, sodium cyanoborohydride.

When -X-X- is -CH=CH-, the compounds of the formula (I) may be prepared using the Wittig or Horner-Emmons reactions. The corresponding compound in which -X-X- is -CH₂CH₂- can be formed by reduction of the -CH=CH- group, for example with hydrogen over a palladium-on-carbon catalyst.


An -X-X- bond of the formula -COO- or -OC(O)- may be formed by reacting the appropriate hydroxy and activated carboxylic acid (e.g. acid chloride or reactive ester) intermediates under conditions known for ester bond formation. Alternatively, a hydroxy and a carboxylic acid intermediate could be reacted together in the presence of diethylazodicarboxylate/triphenylphosphine.

An -X-X- bond of the formula -CH₂O- or -OCH₂- may be formed by reacting the appropriate hydroxy intermediate with

the appropriate alkyl halide in the presence of a base. Conditions for the formation of an ether bond are known in the art.

These reactions can also be used to form intermediates, which contain one of the above -X-X- bonds.

Compounds of the formula (I) may also be prepared by introducing the Lp(D)_n group into a compound of the formula (11):




10

Reactive groups in Lp(D)_n, which could cause side-reactions can of course be protected. The reaction between the compound (11) and H-Lp(D)_n is conveniently carried out in an inert organic solvent, in the presence of an organic base such as an amine (e.g. ethyldiisopropylamine), additionally in the presence of a reagent such as diethylcyanophosphonate.


Intermediates which already contain the Lp(D)_n group may be prepared from the appropriate carboxy compound using similar reaction conditions to those described above.

Compounds of the formula (I) can also be prepared by reacting a compound of the formula (12):



25

or (13):



(wherein X⁵ is a leaving group such as tosyloxy, mesyloxy or


halo) with a compound of formula HNR_aR_b . The reaction between the compound of formula (12) and HNR_aR_b may be carried out in the presence of a reducing agent, such as sodium cyanoborohydride, for example using conditions similar to those described in Examples 1 to 5. The reaction between the compound of formula (13) and HNR_aR_b may be carried out in an inert organic solvent such as THF, in the presence of an inorganic base such as potassium carbonate, and preferably in the presence of sodium iodide. The reaction usually takes place at or near reflux of THF.

Intermediates containing the Lp(D)_n group can also be formed using these reactions from appropriate intermediates, although normally the introduction of the HNR_aR_b group is the last step in the synthesis.

- Hence the present invention also provides a process for the preparation of a compound of formula (I) comprising:
- when $-\text{X}-\text{X}$ is $-\text{CONH}-$, reacting a compound of formula (10) with a compound of formula R_2-COOH , under amide bond-forming conditions;
 - reacting a compound of formula (11) with a compound of formula H-Lp(D)_n under amide bond-forming conditions; or
 - reacting a compound of formula (12) or (13) with a compound of formula HNR_aR_b ;

wherein R_2 , R_a , R_b and Lp(D)_n are as hereinabove defined and formulae (10), (11), (12) and (13) are as hereinabove defined.

A compound of the formula (12) or (13) may be formed from the corresponding hydroxymethyl compound of formula (14).




Oxidation of the hydroxymethyl compound with a suitable mild oxidising agent for the formation of aldehydes, such as *N*-methylmorpholine oxide in the presence of tetrapropylammonium perruthonate, can be used to form a

compound of the formula (12).

Other possible oxidising agents include manganese dioxide or DMSO/oxalyl chloride or DMSO/SO₃ or Dess-Martin reagent.


A compound of formula (13) can be formed from the
5 hydroxymethyl compound of formula (14) by introducing the leaving group X. When X is mesyloxy or tosyloxy, the hydroxymethyl compound may be reacted with the mesyl or tosyl halide in the presence of an organic base, such as triethylamine, in an inert organic solvent such as
10 dichloromethane.

A hydroxymethyl compound of formula (14) in which X-X is CONH may be prepared by reacting a compound of formula (15).




with a compound of formula R₂COOH under amide bond forming
15 conditions, as described previously.


A compound of formula (15) may be prepared by reacting a compound of formula (16).



in which PG is an amino protecting group, such as t-
20 butoxycarbonyl, with a compound of formula (17)



under amide bond forming conditions, to afford a compound of formula (18)




25 followed by removal of the protecting group, PG.


A compound of formula (10) may be prepared by reacting a compound of formula (16) with a compound of formula H-Lp(D)_n under amide bond-forming conditions to afford a compound of

formula (19), followed by removing the protecting group PG.

A compound of formula (11) in which X-X is CONH may be prepared by reacting a compound of formula (20)




- 5 in which PG' represents a carboxyl protecting group with a compound of formula R_2COOH under amide bond-forming conditions to afford a compound of formula (21)



followed by removing the protecting group PG'.

- 10 When -X-X is -CONH- and Y is CH, a compound of formula (I) may be prepared by a number of steps from an amino acid derivative using the reactions described above. For example, see Scheme I.

Scheme 1



PG is an amino-protecting group

PG' is a carboxy-protecting group

An amino acid from Scheme I may be prepared (for example) by one or more of the following methods:

- 5 (i) from aryl or heteroaryl aldehydes via the Strecker synthesis or modifications thereof, via Bucherer-Bergs hydantoin synthesis, or via the Ugi methodology ("Isonitrile

Chemistry", Ugi I. Ed.; Academic: New York, 1971;145-1999, "Multicomponent Reactions with Isocyanides", Domling, A.; Ugi, I. *Angew. Chem. Int. Ed.* 2000, 39, 3168; "Amino Acid Derivatives by Multicomponent Reactions", Dyker, G. *Angew. Chem. Int. Ed. Engl.* 1997, 36, 1700; and also see "A new Class of Convertible Isocyanides in the Ugi Four-Component Reaction", Lindhorst, T.; Bock H.; Ugi, I. *Tetrahedron*, 1999, 55, 7411.) with removal and replacement of protecting groups;

(ii) from styrenes via Sharpless methodology (*J. Am. Chem.*

10 *Soc.* 1998,120, 1207-1217)


(iii) from aryl boronic acids via Petasis methodology (*Tetrahedron*, 1997, 53, 16463-16470) with removal and replacement of protecting groups;

(iv) from aryl and heteroaryl acetic acids - via Evan's
15 azidation (*Synthesis*, 1997, 536-540) or by oximation, followed by reduction and addition of protecting groups; or

(v) from existing aryl glycines by manipulation of functional groups, for example, alkylation of hydroxy groups, palladium assisted carbonylation of triflates derived from hydroxy
20 groups and further manipulation of the carboxylic esters to give carboxylic acids by hydrolysis, carboxamides by activation of the carboxylic acid and coupling with amines, amines via Curtius reaction on the carboxylic acid, or alkylsulphonyl compounds by oxidation of alkylthio compounds;

25 (vi) from aliphatic, carbocyclic and non-aromatic heterocyclic aldehydes and ketones using a Horner-Emmons reaction with N-benzyloxycarbonyl)- α -phosphonoglycine trimethyl ester (*Synthesis*, 1992, 487-490), or by any other method known in the art or

30 (vii) from oximes of formula



in which Pg is a carboxy protecting group, by reduction.

A starting material for the preparation of a compound of formula (I), where the alpha atom is nitrogen, may be

5 produced, for example, by reaction of a beta protected hydrazine (such protection to be chosen as to be compatible with the subsequent reagents to be employed) with phosgene, diphosgene, triphosgene or N,N'carbonyl diimidazole to give a reactive compound of the type
10 $\text{PGNHN}(\text{Cy})\text{COCl}$ or $\text{PGNHN}(\text{Cy})\text{CO-imidazole}$ (wherein PG is a protecting group).

This intermediate may be used as has been described above for the carboxylic starting reagents where the alpha atom is carbon.

15 The skilled person will be aware that at certain stages in the synthesis of a compound of formula (I) it may be necessary to protect a reactive functional group in the molecule to prevent unwanted side-reactions.


The protection of amino and carboxylic acid groups is
20 described in McOmie, Protecting Groups in Organic Chemistry, Plenum Press, NY, 1973, and Greene and Wuts, Protecting Groups in Organic Synthesis, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of carboxy protecting groups include $\text{C}_1\text{-C}_6$ alkyl groups such as methyl, ethyl, t-butyl and t-amyl; aryl($\text{C}_1\text{-}$
25 C_4)alkyl groups such as benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, benzhydryl and trityl; silyl groups such as trimethylsilyl and t-butyl
30 (trimethylsilylmethyl)prop-1-en-3-yl.

Examples of amine protecting groups (PG) include acyl

groups, such as groups of formula RCO in which R represents C₁₋₆ alkyl, C₃₋₁₀ cycloalkyl, phenyl C₁₋₆ alkyl, phenyl, C₁₋₆ alkoxy, phenyl C₁₋₆ alkoxy, or a C₃₋₁₀ cycloalkoxy, wherein a phenyl group may be optionally substituted, for example by one or two
5 of halogen, C₁₋₄ alkyl and C₁₋₄ alkoxy.

Preferred amino protecting groups include benzyloxycarbonyl (CBz), t-butoxycarbonyl (Boc) and benzyl.

In another aspect the invention relates to a process for preparing a compound of formula I comprising deprotecting a
10 compound of formula (I'):



wherein R^{2'} is R² (as hereinabove defined) or protected R², Cy' is Cy (as hereinabove defined) or protected Cy and Lp(D)_n' is Lp(D)_n (as hereinabove defined) or protected Lp(D)_n; providing
15 at least one protecting group is present.

If necessary physiologically tolerable salts can be formed using methods known in the art.

It will be understood that the compounds of formula (I) may be isolated in the form of salts or solvates (which may or
20 may not be physiologically tolerable), and that all such salts and solvates are therefore included within the scope of the present invention.

All novel intermediates described herein are provided as further aspects of the invention.

25 The compounds of the invention may be administered by any convenient route, e.g. into the gastrointestinal tract (e.g. rectally or orally), the nose, lungs, musculature or vasculature or transdermally. The compounds may be administered in any convenient administrative form, e.g.
30 tablets, powders, capsules, solutions, dispersions, suspensions, syrups, sprays, suppositories, gels, emulsions, patches etc. Such compositions may contain components

conventional in pharmaceutical preparations, e.g. diluents, carriers, pH modifiers, sweeteners, bulking agents, and further active agents. Preferably the compositions will be sterile and in a solution or suspension form suitable for
5 injection or infusion. Such compositions form a further aspect of the invention.

The following are examples of pharmaceutical compositions of compounds according to the invention.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

5	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
10	Magnesium stearate <u>10</u>
	Total 460 mg

The above ingredients are mixed and filled into hard
15 gelatin capsules in 460 mg quantities.

Formulation 2

Tablets each containing 60 mg of active ingredient are made as follows:

5

Active Ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
10 Polyvinylpyrrolidone	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
15 Total	150 mg

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders which are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Viewed from this aspect the invention provides a pharmaceutical composition comprising a serine protease inhibitor according to the invention together with at least one pharmaceutically acceptable carrier or excipient. The pharmaceutical composition may also optionally comprise at least one further antithrombotic and/or thrombolytic agent.

Viewed from a further aspect the invention provides the use of a serine protease inhibitor according to the invention for the manufacture of a medicament for use in a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat (i.e. treat or prevent) a condition responsive to said inhibitor.

Viewed from a further aspect the invention provides a method of treatment of the human or non-human animal body (e.g. a mammalian, avian or reptilian body) to combat a condition responsive to a serine protease inhibitor (e.g. a condition such as a thrombotic disorder responsive to a factor Xa inhibitor), said method comprising administering to said body an effective amount of a serine protease inhibitor according to the invention.

The dosage of the inhibitor compound of the invention will depend upon the nature and severity of the condition being treated, the administration route and the size and species of the patient. However in general, quantities of from 0.01 to 100 $\mu\text{mol/kg}$ bodyweight will be administered.

All publications referred to herein are hereby incorporated by reference.

The invention will now be described further with reference to the following non-limiting Examples.

Experimental

Abbreviations used follow IUPAC-IUB nomenclature.

- 5 Additional abbreviations are HPLC, high-performance liquid chromatography; rPHPLC, reverse phase HPLC; SCX, strong cation exchange resin; THF, tetrahydrofuran; HOAc, acetic acid; DMSO, dimethyl sulfoxide (perdeuterated if for NMR); EtOAc, ethyl acetate; EtOH, ethanol; DMF, dimethylformamide; DCM, 10 dichloromethane; HOAT, 1-hydroxy-7-azabenzotriazole; HOBT, 1-hydroxy benzotriazole; EDCI, 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; DIPEA, diisopropylethylamine; Boc, tertiary butyloxycarbonyl; TEA, triethylamine; TFA, trifluoroacetic acid; MALDI-TOF, Matrix assisted laser 15 desorption ionisation-time of flight mass spectrometry, CI-MS, chemical ionization mass spectrum; API-MS (atmospheric pressure chemical ionization mass spectra) were obtained on a PE Sciex (trademark) API 150EX with a heated nebulizer and nitrogen as the reagent gas in positive ion mode. RT, 20 retention time; TLC, thin layer chromatography with R_f as relative mobility. All solution concentrations are expressed as %Vol./%Vol. unless otherwise stated. Reagents were obtained from a variety of commercial sources.
- 25 IR means an infrared spectrum was obtained. ^1NMR , $^1\text{H-NMR}$, or $^1\text{H NMR}$ means a proton magnetic resonance spectrum was obtained.

- In general in this specification, "D-" or "R-" in the 30 name of a product indicates the product was made beginning with a chiral starting material, for example D-phenylglycine; however, racemization may have occurred, and the enantiomeric purity may not have been determined.

General Experimental Procedures:**Purification of Compounds (rpHPLC chromatography)**

Material is or was purified using standard reverse-phase preparative chromatography techniques. A 5 micron, 20 x 50 mm 5 O.D. C18 column is or was used (YMC ODS-A) with a flow rate of 20 mL/min and an standard elution time of 10-15 minutes. A gradient of water:acetonitrile (between 95:5 to 5:95; each eluent w/0.1% TFA) over the elution time is or was used. Fractions containing product are or were concentrated, frozen, 10 and lyophilized to afford, when applicable, the trifluoroacetate salt of the product. The free base can or could be obtained, if desired, by loading a methanolic solution of the trifluoroacetate salt onto an ion-exchange resin (SCX, Varian) and subsequent elution of the resin with 15 methanol followed by 2 N ammonia in methanol. Concentration of the later fractions affords or afforded the free base product. Preparation of a hydrochloride salt from the free base is or was completed by treatment an organic solution of the free base (EtOAC, methylene chloride) with anhydrous HCl 20 in diethyl ether and concentration.

Preparation of Starting Materials and Intermediates

Intermediate substituted glycine compounds for starting materials and intermediates, including those in which the 25 amino group and/or the carboxy group is protected, conveniently may be prepared using one of the procedures below, or by a similar procedure. It may be convenient or preferred to change the order of steps in the preparation of a compound of the invention and to use a similar procedure with 30 a different intermediate. In particular, it may be convenient to use an acyl group R₂-CO- initially in a preparation, rather than an amino protecting group.

Abbreviations, in addition to others listed herein,

include: TEMPO: 2,2,6,6-tetramethyl-1-piperidinyloxy, free radical; (DHQD)₂PHAL: hydroquinidine 1,4-phthalazinediyl diether; r.b. or rb, round bottomed; PPh₃, triphenylphosphine; Boc₂O or Boc anhydride: di-tert-butyl dicarbonate.

5

Preparation of Intermediates KE-1 - KE-5

The following compounds were prepared according to the indicated method (Method KE-A) from the indicated starting materials, unless otherwise described.

10

Intermediate KE-1

Ethyl oxo-quinolin-8-ylacetate.

Method KE-A

To a stirring solution of 8-bromoquinoline (10.1 g, 48.5 mmol) in THF (500 mL) at -78 °C was added dropwise a 1.3 M solution of sec-butyl lithium (37.3 mL, 48.5 mmol) in cyclohexane. After 5 min, diethyl oxalate (8 mL, 58.3 mmol) was added; and the solution was allowed to slowly warm to room temperature overnight. The next morning, the reaction was quenched with the addition of saturated aqueous NH₄Cl; and the solvent was removed in vacuo. The residue was partitioned between ethyl acetate and satd aq. NaHCO₃; the layers were separated; and then the aqueous phase was washed with brine, dried with MgSO₄, filtered and concentrated in vacuo. The residue was chromatographed over silica gel, eluting with 20% ethyl acetate/hexanes through 25% ethyl acetate/hexanes. The product containing fractions were combined and concentrated in vacuo to give 5.88 g (53%) of the title compound.

30 1H-NMR

IS-MS, m/e 230.1 (M+1)

Intermediate KE-2

Ethyl oxo-quinolin-5-ylacetate.

Prepared from 5-bromoquinoline and diethyl oxalate using Method KE-A.

¹H-NMR

5 IS-MS, m/e 230.0 (M+1)

Intermediate KE-3

Ethyl oxo-thiazol-5-ylacetate.

To a r.b. flask (500 cm³) under argon, fitted with
10 ethanol thermometer, septum cap, and dropping funnel, was added anhydrous ether (100 cm³) with stirring. This was cooled to -78 °C and 2 M n-butyllithium (60 cm³, 120 mmol) was added.

A solution of silyl thiazole (16 g, 16 cm³, 100 mmol) in anhydrous ether (100 cm³) was then added by dropping funnel
15 over 30 minutes. This was allowed to stir for 1 hour to give a peach suspension. To this was added diethyl oxalate (16.3 cm³, 17.5 g, 120 mmol) rapidly to give a brown solution, resulting in a temperature increase to -30 °C. This was allowed to cool back to -78 °C and stirred for 30 minutes.

20 Reaction monitored by ¹H NMR (CDCl₃).

The brown solution was poured onto 5% hydrochloric acid solution (300 cm³) with vigorous stirring for 30 minutes. Ether layer was separated and washed with saturated bicarbonate (ca. 80 cm³), dried over magnesium sulphate, and
25 concentrated in vacuo to give an orange oil. This was purified by flash chromatography (10% ethyl acetate/hexane) to give a yellow oil (7.31 g, 39.47 mmol) [40% Yield].

¹H NMR (CDCl₃); 1.42 (3H, t), 4.45 (2H, q), 8.89 (1H, s), 9.10
30 (1H, s).

Intermediate KE-4

Ethyl oxo-thiazol-2-ylacetate.

Prepared from thiazole and diethyl oxalate using Method KE-A. In this case the temperature was held at -35 °C and n-butyllithium in hexane was used in place of sec-butyllithium in cyclohexane.

5 ¹NMR

IS-MS, m/e 165.0 (M+1)

Intermediate KE-5

Ethyl oxo-isoquinolin-8-ylacetate.

10 Prepared from 8-bromoisoquinoline and diethyl oxalate using Method KE-A, substituting n-butyl lithium in hexanes for sec-butyl lithium in cyclohexane.

¹NMR

IS-MS, m/e 230.0 (M+1)

15 Analysis for C₁₃H₁₁NO₃:

Calcd: C, 68.11; H, 4.84; N, 6.11;

Found: C, 68.11; H, 5.00; N, 6.14.

Preparation of Intermediates OX-1 - OX-9

20 The following compounds were prepared according to the indicated method (Method OX-A or Method OX-B) from the indicated starting materials unless otherwise described.

Intermediate OX-1

25 Ethyl Hydroxyimino-pyridin-2-ylacetate.

Method OX-A

To a stirring solution of ethyl 2-pyridylacetate (12.6 g, 76.3 mmol) in acetic acid (19 mL) at 5 °C was added a solution of sodium nitrite (6.05 g, 87.7 mmol) in water (12 mL) at a
30 rate sufficient to maintain the internal temperature below 15 °C. After complete addition and an additional 30 min, an additional 30 mL of water were added. The resulting white precipitate was filtered, washed with water, satd aq. NaHCO₃, and again with water. The solid was then dried under vacuum

to give 14.1 g (95%) of the title compound.

¹H-NMR

IS-MS, m/e 194.9 (M+1)

5 Analysis for C₉H₁₀N₂O₃:

Calcd: C, 55.67; H, 5.19; N, 14.43;

Found: C, 55.79; H, 5.14; N, 14.13.

Intermediate OX-2

10 Ethyl Hydroxyimino-pyridin-3-ylacetate.

Using the procedure of Tikk et al [Acta. Chimica, Hungarica, 114(3-4), 355], a mixture of ethyl hydroxyimino-pyridin-3-yl-acetate and n-butyl hydroxyimino-pyridin-3-yl-acetate was prepared from ethyl 3-pyridinylacetate and n-butyl
15 nitrite.

¹H-NMR

IS-MS, m/e 195 (M+1), 223.1 (M+1)

20 Intermediate OX-3

Ethyl Hydroxyimino-quinolin-8-ylacetate.

Method OX-B

To a stirring solution of ethyl oxo-quinolin-8-yl-acetate (5.5 g, 24 mmol) in ethanol (140 mL) was added sodium acetate
25 (2.16 g, 26.4 mmol) followed by hydroxylamine hydrochloride (2.67 g, 38.4 mmol). The mixture was heated to reflux; and, after 7 h, the heating mantle was removed and the solution was allowed to stir overnight at room temperature. The next
30 morning, the solvent was removed in vacuo and the residue was partitioned between ethyl acetate and satd aq. NaHCO₃. The layers were separated and the organic phase was washed with brine, dried with Na₂SO₄, filtered and concentrated in vacuo.
The resulting foam was recrystallized from dichloromethane/hexanes to give an initial crop of 2.5 g of

the title compound as an off-white solid, followed by 0.31 g of a second crop. The mother liquor was then concentrated in vacuo, the residue was dissolved in a minimal amount of dichloromethane. The solution was then chromatographed over silica gel, eluting with 30% ethyl acetate/hexanes, then 40% ethyl acetate/hexanes, and finally with ethyl acetate. The product containing fractions were combined and concentrated in vacuo to give 1.94 g of the title compound for a combined yield of 4.75 g (81%).

10

1H-NMR

IS-MS, m/e 245.0 (M+1)

Intermediate OX-415 **Ethyl Hydroxyimino-quinolin-5-ylacetate.**

Prepared from ethyl oxo-quinolin-5-yl-acetate using Method OX-B.

1H-NMR

20 IS-MS, m/e 245.0 (M+1)

Intermediate OX-5**Ethyl Hydroxyimino-thiazol-5-ylacetate.**

To a r.b. flask (500 cm³) was added the ethyl oxo-thiazol-5-ylacetate (6.30g, 34.02 mmol) to ethanol (ca. 180 cm³) with stirring. Sodium acetate (3.06g, 37.30 mmol) and hydroxylamine hydrochloride (3.78g, 54.43 mmol) were then added to give an off-white suspension. This was brought to reflux at 85 °C for 1 hour. Reaction monitored by TLC (60% hexane/ethyl acetate; s.m. r.f 0.5, prod. r.f. 0.3.). Reaction cooled and concentrated in vacuo. Product taken up in ethyl acetate (c.a. 200 cm³) and washed with 5% hydrochloric acid solution. Ethyl acetate layer was dried over magnesium sulphate and evaporated to dryness to give a

cream solid (6.372g, 31.825 mmol) [94% Yield].

^1H NMR (CDCl_3); 1.40 (3H, m), 4.40 (2H, m), 8.06 ($1/3\text{H}$, s), 8.78 ($1/3\text{H}$, s), 8.95 ($2/3\text{H}$, s), 8.98 ($2/3\text{H}$, s).

5

Intermediate OX-6

Ethyl α -Oximino-thiazole-4-acetate.

To a 2 necked r.b. flask (100 cm^3) with ethanol thermometer, concentrated sulphuric acid (25 cm^3) was added
10 and cooled to 0 $^\circ\text{C}$ with stirring. To this solution was added the ethyl α -oximino-2-aminothiazole-4-acetate (5.00 g, 23.231 mmol). Water (10 cm^3) was then added and cooled to -10 $^\circ\text{C}$. A solution of sodium nitrite (1.683 g, 24.393 mmol) in water (5 cm^3) was then added slowly over an hour keeping the
15 temperature below -5 $^\circ\text{C}$.

To a separate r.b. flask (500 cm^3), water (180 cm^3) was added and cooled to 3 $^\circ\text{C}$. The reaction solution was poured in to the cold water with stirring and then cooled to -5 $^\circ\text{C}$. To this solution, 50% hypophosphoric acid (90 cm^3) was added
20 dropwise over 10 minutes keeping the temperature at -5 $^\circ\text{C}$. The solution was allowed to warm to room temperature and stirred overnight. The product was extracted with diethyl ether (ca. 3x150 cm^3) and washed with water. The ether layer was concentrated in vacuo and treated to flash chromatography (50%
25 ethyl acetate/n-hexane) to yield a orange oil upon concentration in vacuo (0.60 g, 3.00 mmol) [13% yield].

^1H NMR (CDCl_3) 1.35 (3H, m), 4.35 (2H, m), 8.4 (1H, s), 8.9 (1H, s), 14.4 (1H, s).

30

Intermediate OX-7

Ethyl α -Oximino-2-methylthiazole-4-acetate.

This was prepared from ethyl- γ -chloro- α -oximino-

acetoacetate (1.44g) using the method of Hatanaka et al.
(*Journal of Medicinal Chemistry*, 1973, 16(9), 978-984) to
yield the titled compound (0.64 g).

5 ^1H NMR (CDCl_3) 1.35 (3H, t), 2.7 (3H, s), 4.35 (2H, q), 8.2
(1H, s).

Ethyl γ -Chloro- α -oximinoacetoacetate.

This was prepared from ethyl oximinoacetoacetate (1.73 g)
10 using the method of Hatanaka et al. (*Journal of Medicinal
Chemistry*, 1973, 16(9), 978-984) to yield the titled compound
(1.44g).

^1H NMR (CDCl_3) 1.25 (3H, t), 4.3 (2H, q), 4.55 (2H, s), 9.45
15 (1H, s), contains 20% starting material by NMR.

Ethyl Oximinoacetoacetate

This was prepared from ethyl acetoacetate (10.00g) using
the method of Fischer (*Organic Synthesis Coll. Vol. 3*, 513-
20 516) to yield the titled compound (12.45 g).

^1H NMR (CDCl_3) 1.25 (3H, t), 2.35 (3H, s), 4.3 (2H, q), 8.8
(1H, br.).

25 Intermediate OX-8

Ethyl hydroxyimino-thiazol-2-ylacetate.

Prepared from ethyl oxo-thiazol-2-ylacetate using Method
OX-B.

30 ^1NMR

IS-MS, m/e 198.9(M-1)

Intermediate OX-9**Ethyl hydroxyimino-isoquinolin-8-ylacetate.**

Prepared from ethyl oxo-isoquinolin-8-ylacetate using Method OX-B.

5

¹H NMR

IS-MS, m/e 245.0 (M+1)

Analysis for C₁₃H₁₂N₂O₃:

Calcd: C, 63.93; H, 4.95; N, 11.47;

10 Found: C, 63.68; H, 4.60; N, 11.34.

Preparation of Intermediates AL-1 - AL-3

The following compounds were prepared according to the indicated method (Method AL-A or Method AL-B) from the indicated starting materials, unless otherwise described.

Intermediate AL-1**R-3-Bromo-(1-t-butoxycarbonylamino-2-hydroxyethyl)benzene.****Method AL-A**

20 Sodium hydroxide (3.33 g, 83.25 mmol) was dissolved in water (220 mL), and 20 mL of the resulting solution was removed and added to potassium osmate (410 mg, 1.11 mmol). The remaining sodium hydroxide solution (200 mL) was added to a stirred solution of t-butyl carbamate (9.9 g, 84.5 mmol) in 25 n-propanol (110 mL) followed by freshly prepared t-butyl hypochlorite (9.65 mL; 83.5 mmol). After stirring for 5 min, the solution was cooled to 0 °C. A solution of (DHQD)₂PHAL (1.30 g, 1.67 mmol) in n-propanol (110 mL) was added, followed by a solution of 3-bromostyrene (5 g, 27.31 mmol) in n- 30 propanol (220 mL), followed by dropwise addition of the potassium osmate/sodium hydroxide solution. The reaction was stirred overnight. Saturated aqueous sodium sulfite (150 mL) was added, and the reaction was stirred for 15 min. The aqueous layer was separated and extracted with ethyl acetate

(3x 200 mL). The combined organic layers were washed with brine and dried over MgSO_4 . Removal of solvent under vacuum gave the crude product which was purified by chromatography (silica, 3:2 hexane:ethyl acetate then rechromatographed loading with toluene, gradient elution with hexane - 4:1 hexane:ethyl acetate) to give the title product (4.18 g, 49%).

Melting Point = 90-91 °C

^1H NMR (CDCl_3).

Intermediate AL-2

R-3-Methoxycarbonyl-(1-t-butoxycarbonylamino-2-hydroxy-ethyl)benzene.

Method AL-B

In a glass liner containing a stirrer bar was placed $\text{Pd}(\text{OAc})_2$ (871 mg, 3.88 mmol), PPh_3 (1.96 g, 7.47 mmol), NaOAc (1.48 g, 18.04 mmol) and DMF (82 mL). To this stirred solution was added a solution of R-3-bromo-(1-t-butoxycarbonylamino-2-hydroxyethyl)benzene (4.27 g, 13.5 mmol) in MeOH (82 mL). The resulting solution was purged with nitrogen and placed in a stirred pressure vessel. The system was charged to 4.1 bar (60 psig) of CO and heated at 95 °C for 36 h. The mixture was cooled to room temperature, filtered through diatomaceous earth, and partitioned between ethyl acetate and water. The organic layer was washed with water (3x) and brine (1x) and dried over MgSO_4 . Removal of solvent under vacuum gave the crude product which was purified by chromatography (silica gel, gradient elution with 30-35% ethyl acetate/hexane) to provide the title product (3.53 g, 89%).

Melting Point = 73-75 °C with decomposition

^1H NMR (CDCl_3).

API-MS, m/e = 240 ($\text{M}-\text{C}_4\text{H}_9+1$).

Intermediate AL-3**R-3-Cyano- (1-t-butoxycarbonylamino-2-hydroxyethyl)benzene.**

Prepared from 3-cyanostyrene using Method AL-A.

3-Cyanostyrene was prepared using the method described below.

5

Melting Point = 76 °C.

¹H NMR (CDCl₃).

Preparation of 3-Cyanostyrene.

- 10 To a stirred suspension of methyltriphenylphosphonium
bromide (75 g, 209.71 mmol) in dry THF (750 mL) at 0 °C under
nitrogen was added dropwise n-BuLi (83 mL, 2.5 M in hexanes,
207.50 mmol). The mixture was warmed to room temperature. 3-
Cyanobenzaldehyde (25 g, 190.65 mmol) was added as a solid in
15 5 g batches, and the mixture was stirred at room temperature
overnight. The reaction was quenched in water, and the
solvent was removed under vacuum. The residue was dissolved
in the minimal amount of THF, and triphenylphosphine oxide was
precipitated using ether. The solid was filtered through
20 diatomaceous earth, and the filtrate was concentrated.
Distillation by Kugelrohr at 90 °C/33 Pa (0.25 mm Hg) gave the
product as a colorless oil (15.5 g, 62%).

Boiling Point = 90 °C at 0.25 mmHg.

25 ¹H NMR (CDCl₃).

Preparation of Intermediates PAE-1 - PAE-18

- The following compounds were prepared according to the
indicated method (Method PAE-A, Method PAE-B, Method PAE-C,
30 Method PAE-D or PAE-E) from the indicated starting materials,
unless otherwise described.

Intermediate PAE-1**Boc-D,L-(2-pyridinyl)glycine Ethyl Ester.****Method PAE-A**

To a solution of ethyl hydroxyimino-pyridin-2-yl-acetate
5 (7.8 g, 40.15 g) in ethanol (175 mL) and glacial acetic acid
(20 mL) was added 5% Pd/C, and the mixture was shaken in a
hydrogenation apparatus under an atmosphere of hydrogen at 4.1
bar (45 psig) for 4 h. The mixture was filtered through
diatomaceous earth and concentrated in vacuo. The residue was
10 dissolved in THF/H₂O (1/1, 240 mL) and treated with di-tert-
butyl dicarbonate (14.23 g, 65.2 mmol) and sodium bicarbonate
(27.4 g, 326 mmol). After stirring at room temperature for
2 h, the solution was concentrated in vacuo and the residue
was partitioned between EtOAc and water. The organic phase
15 was washed with brine, dried over magnesium sulfate, filtered
and concentrated in vacuo. The crude material was purified
via chromatography over silica gel, eluting with a stepwise
gradient of 10-20% ethyl acetate in dichloromethane to give
8.11 g (72%) of the title compound as a yellow oil.

20

1H-NMR

IS-MS, m/e 281.1 (M+1)

Intermediate PAE-2**25 Boc-D,L-(3-pyridinyl)glycine Ethyl Ester.**

Prepared from ethyl hydroxyimino-pyridin-3-ylacetate
using Method PAE-A.

1H-NMR

30 IS-MS, m/e 281.1 (M+1)

Intermediate PAE-3**Boc-D,L-(8-quinolinyl)glycine Ethyl Ester.****Method PAE-B**

To a stirring solution of ethyl hydroxyimino-quinolin-8-ylacetate (2.4 g, 9.8 mmol) in 50% aq. formic acid (50 mL) at 0 °C was added zinc dust (2 g, 31 mmol). After 1 min, the mixture was filtered through diatomaceous earth and the filtrate was loaded onto an SCX column. After washing the column with methanol, the product was eluted with a 3 to 1 mixture of dichloromethane and (2 N NH₃ in methanol). The product containing fractions were combined and concentrated in vacuo to give 2.24 g of light orange oil (IS-MS, m/e 231.0 (M+1)).

The oil (2.14 g, 9.3 mmol) was dissolved in THF (40 mL) and to this stirring solution was added triethylamine (1.4 mL, 10.2 mmol), followed by di-tert-butyl dicarbonate (2.1 g, 9.8 mmol). After 45 min, the solvent was removed in vacuo and the residue was partitioned between ethyl acetate and water. The organic phase was then washed with satd aq. NaHCO₃, dried with Na₂SO₄, filtered and concentrated in vacuo. The residue was dissolved in a minimum volume of dichloromethane and chromatographed over silica gel, eluting with 5% ethyl acetate in hexanes. The product containing fractions were combined and concentrated to give 2.5 g (81%) of the title compound.

1H-NMR

IS-MS, m/e 331.0 (M+1)

Intermediate PAE-4**Boc-D,L-(5-quinolinyl)glycine Ethyl Ester**

Prepared from ethyl hydroxyimino-quinolin-5-ylacetate using Method PAE-B.

1H-NMR

IS-MS, m/e 331.0 (M+1)

Intermediate PAE-5

N-4-Methoxybenzoyl-N-2,4-dimethoxybenzyl-D,L-(2-trifluoro-
5 methylphenyl)glycine Methyl Ester.

Method PAE-C

To 2-trifluoromethylbenzaldehyde (1 g, 5.7 mmol) with stirring was added 2,4-dimethoxybenzylamine (0.86 mL, 5.7 mmol) and methanol (2 mL). After 5 min, the solution was
10 diluted with toluene 100 mL and concentrated in vacuo (twice).

The residue was then dissolved in anhydrous methanol (12 mL) and 1,1-dimethyl-2-(methoxycarbonyloxy)ethyl isonitrile [Tetrahedron, 55 (1999) 7411-7420] (0.9 g, 5.7 mmol) was added, followed by 4-methoxybenzoic acid (0.87 g, 5.7 mmol).

15 After stirring for 72 h, the solvent was removed in vacuo and the residue was chromatographed over silica gel, eluting with a step gradient of 30% ethyl acetate in hexanes through 50% ethyl acetate in hexanes. The product containing fractions were combined and concentrated in vacuo; and then the residue

20 was dissolved in ethyl acetate, washed with satd aq. NaHCO₃, dried with Na₂SO₄, filtered and concentrated to give 1.76 g (48%) of thick oil (NMR, IS-MS, m/e 633.0 (M+1)). The oil (0.5 g, 0.79 mmol) was then dissolved in toluene (5 mL) and concentrated in vacuo (twice) to give a white foam. The

25 residue was then dissolved in THF (3 mL) and potassium tert-butoxide (0.11 g, 0.95 mmol) was added. After 15 min, 12 N HCl (0.079 mL, 0.95 mmol) was added and the solution was allowed to stand overnight in the refrigerator. The next morning, the solvent was removed and the residue was

30 chromatographed over silica gel, eluting with 30% ethyl acetate in hexanes. The product containing fractions were combined and concentrated to give 0.32 g (79%) of the title compound.

¹H-NMR

IS-MS, m/e 518.0 (M+1)

Intermediate PAE-6

5 BOC-D,L-(5-thiazolyl)glycine ethyl ester.

To a r.b. flask (250 cm³), D,L-(5-thiazolyl)glycine ethyl ester (4.60 g, 24.7 mmol) was added to tetrahydrofuran (c.a. 100 cm³) with stirring to give a yellow solution. BOC anhydride (5.439g, 24.948 mmol) and triethyl amine (3.79 cm³, 10 2.75g, 27.17 mmol) were then added with stirring for 1 hour. Reaction monitored by TLC (60% hexane/ethyl acetate; s.m. r.f 0.05, prod. r.f. 0.5.). The reaction concentrated in vacuo and product taken up in ethyl acetate (c.a. 150 cm³), washed with 5% hydrochloric acid solution (c.a. 30 cm³), and 15 saturated bicarbonate (ca. 30 cm³). Ethyl acetate layer was dried over magnesium sulphate and evaporated to dryness to give an orange oil (7.42 g, ~24.70 mmol) [-100% Yield].

¹H NMR (CDCl₃); 1.30 (3H, t), 1.48 (9H, s), 4.28 (2H, q), 5.68 20 (1H, br.), 7.88 (1H, s), 8.78 (1H, s).

D,L-(5-Thiazolyl)glycine Ethyl Ester.

To a r.b. flask (250 cm³), was added 5-thiazolyl-oximinoacetic acid ethyl ester (6.37 g, 31.825 mmol) to 25 ethanol (c.a. 80 cm³) with stirring. 50% Formic acid solution (50 cm³) was added with zinc dust (5.10 g, 81.83 mmol) and allowed to stir overnight. Reaction monitored by TLC (60% hexane/ethyl acetate; s.m. r.f 0.3, prod. r.f. 0.05.). Reaction solution filtered over diatomaceous earth and 30 filtrate concentrated in vacuo. This was basified to pH 9 with anhydrous potassium carbonate and product taken up in 3:1 chloroform/isopropanol solution (c.a. 200 cm³). This was washed with saturated bicarbonate (c.a. 50 cm³), dried over magnesium sulphate and concentrated in vacuo to give a brown

oil (4.60 g, 24.70 mmol) [78% Yield].

^1H NMR (CDCl_3); 1.25 (3H, t), 1.95 (2H, br.), 4.22 (2H, q), 4.85 (1H, s), 7.80 (1H, s), 8.70 (1H, s).

5

Intermediate PAE-7

N-Boc-D,L-(4-thiazolyl)glycine ethyl ester

To a solution of D,L-(4-thiazolyl)glycine ethyl ester (0.460 g, 2.470 mmol) in tetrahydrofuran (20 cm^3), was added

10 di-tert-butyl dicarbonate (0.530 g, 2.470 mmol) and triethylamine (0.344 cm^3 , 2.470 mmol). This was allowed to stir for 1 hour and the solution concentrated in vacuo. The oil was taken up in ethyl acetate (c.a. 50 cm^3) washed with 0.5% hydrochloric acid solution (c.a. 20 cm^3), and saturated
15 sodium bicarbonate solution (c.a. 20 cm^3). This was then dried over magnesium sulphate and concentrated in vacuo to yield an orange oil (0.709 g, 2.477 mmol) [~100% yield].

^1H NMR (CDCl_3) 1.15 (3H, t), 1.35 (9H, s), 4.1 (2H, m), 5.45
20 (1H, d), 5.75 (1H, d), 7.3 (1H, d), 8.7 (1H, d).

D,L-(4-Thiazolyl)glycine Ethyl Ester.

This was prepared from ethyl- α -oximino-thiazole-4-acetate (0.60 g) using the method of Hatanaka et al. (*Journal of*
25 *Medicinal Chemistry*, 1973, 16(9), 978-984) to yield the titled compound (0.46 g).

^1H NMR (CDCl_3) 1.25 (3H, t), 1.8-2.3 (2H, br.), 4.1 (2H, m), 4.75 (1H, s), 7.25 (1H, d), 8.7 (1H, d).

30

Intermediate PAE-8

N-Boc-D,L-(2-methylthiazol-4-yl)glycine Ethyl Ester

To a solution of D,L-(2-methylthiazol-4-yl)glycine ethyl ester (0.397 g, 1.982 mmol) in tetrahydrofuran (20 cm^3), was

added di-tert-butyl dicarbonate (0.475 g, 2.180 mmol) and triethylamine (0.304 cm³, 2.180 mmol). This was allowed to stir for 1 hour and the solution concentrated in vacuo. The oil was taken up in ethyl acetate (c.a. 50 cm³) washed with 5 0.5% hydrochloric acid solution (c.a. 20 cm³), and saturated sodium bicarbonate solution (c.a. 20 cm³). This was then dried over magnesium sulphate and concentrated in vacuo to yield a yellow oil (0.654 g, 2.177 mmol) [-100% yield].

10 ¹H NMR (CDCl₃) 1.1 (3H, s), 1.35 (9H, s), 2.6 (3H, s), 4.15 (3H, m), 5.3 (1H, d), 5.7 (1H, s), 7.0 (1H, s).

D,L-(2-Methylthiazol-4-yl)glycine Ethyl Ester.

This was prepared from ethyl- α -oximino-2-methylthiazole-

15 4-acetate (0.62 g) using the method of Hatanaka et al.
(*Journal of Medicinal Chemistry*, 1973, 16(9), 978-984) to yield the titled compound (0.40 g).

¹H NMR (CDCl₃) 1.15 (3H, t), 1.95 (2H, br.), 2.6 (3H, s), 4.15
20 (2H, m), 4.65 (1H, s), 6.95 (1H, s).

Intermediate PAE-9

Boc-R-(4-Hydroxyphenyl)glycine Methyl Ester

To a stirred mixture of R-(4-hydroxyphenyl)glycine methyl
25 ester hydrochloride (14g) and sodium bicarbonate (11.7 g) in THF (150 mL) and water (50 mL), was added in one portion, di-tert-butyl dicarbonate (15.9 g). The mixture was stirred rapidly to allow thorough mixing for 4 h. Hexane (75 mL) was added and the organic layer separated and washed with satd sodium
30 bicarbonate solution, then brine and then dried with magnesium sulphate. The drying agents was filtered off and washed with a little THF and evaporated to dryness, finishing with a high vacuum pump to remove the last traces of di-t-butyl

dicarbonate. Yield 19.7 g, 96%.

^1H NMR

5 **R-(4-Hydroxyphenyl)glycine Methyl Ester Hydrochloride.**

To a dry 250 mL three necked round bottom flask, equipped with a low temperature thermometer, a septum for nitrogen coverage and another for introduction of thionyl chloride by syringe, was added R-4-hydroxyphenylglycine (12.5 g) and dry
10 methanol (24 mL). The mixture was stirred (magnetic stirrer) and cooled to an internal temperature of -20 °C using cardice/acetone. Using a syringe, thionyl chloride was added dropwise to the cooled mixture over a period of 10 min.

(Care: the reaction of thionyl chloride with methanol is very
15 exothermic and rate of addition should be such that the thionyl chloride is efficiently stirred into the mixture and that the temperature does not rise above -20 °C. Once the addition was complete the mixture was allowed to warm to room temperature overnight (16-18 h). Dry ether (150 mL) was added
20 and the white ppt. that formed was filtered off, washed with a little more ether and dried. Yield 15.5 g, 95%.

^1H NMR

25 **Intermediate PAE-10**

Boc-R-(4-Trifluoromethanesulphonyloxyphenyl)glycine Methyl Ester Hydrochloride.

To a stirred solution of Boc-R-(4-hydroxyphenyl)glycine methyl ester (19 g) in dichloromethane (400 mL) was added 2,6-
30 lutidine (9.44 mL) and 4-dimethylaminopyridine (1.65 g) and the mixture cooled in an ice bath. Trifluoromethanesulphonic anhydride (13.74 mL) was added over a period of 5 min, and then the reaction left to warm to room temperature over 4 h. The organic solution was washed with water (2 x 150

mL), 1 N HCl (2 x 150 mL), and then saturated sodium bicarbonate (150 mL). The organics were dried with magnesium sulphate and then evaporated to an oil. The mixture was purified using flash chromatography (SiO₂ 250 g, eluting with 5 1:1 hexane/dichloromethane and then neat dichloromethane). Pure product fractions were combined and evaporated, finishing with a high vacuum pump to remove all traces of solvent, to give a white solid, 19 g, 77%.

10 ¹H NMR

Intermeidate PAE-11

Boc-R-(4-Methoxycarbonylphenyl)glycine Methyl Ester.

Method PAE-D

15 Boc-R-4-trifluoromethanesulphonyloxyphenylglycine methyl ester (15 g), methanol (32.6 mL), bis-1,3-diphenylphosphinylpropane (448 mg), palladium (II) acetate (255 mg), triethylamine (10.2 mL) and dimethylformamide (72 mL) were placed in the glass liner of pressure (Parr) reactor and the
20 reactor assembled. The vessel was pressurised to ~0.68 bar (10 psig) with nitrogen and the gas released (repeated five times to remove all oxygen from the system). Carbon monoxide gas was then carefully introduced (use extreme care -the gas cylinder is pressurised to far beyond the bursting disc
25 pressure of the Parr, ideally use a pressure regulator to reduce the pressure to ~ 6.8 bar, 100 psig) to ~1.4 bar (20 psig) and released three times (into the back of a fume hood). Carbon monoxide was then added to ~6.8 bar (100 psig) and the stirrer started. The vessel was slowly heated to 65 °C
30 internal temperature and then stirred at 65 °C overnight. (At the early stages more carbon monoxide was added to maintain ~6.8 bar, 100 psig.) A sample was removed after 18 h and examined by tlc. When complete, the reaction was cooled to

-30 °C, the gas released and the vessel flushed five times with nitrogen as before. The reaction mixture was partitioned between ethyl acetate and water, and the organic layer washed with 1 M hydrochloric acid and then saturated sodium bicarbonate. The solution was dried with MgSO_4 and evaporated. Flash chromatography of the resulting oil gave the product, pure by tlc, 10.6 g, 90%.

^1H NMR

10

Intermediate PAE-12

Boc-R-(4-Benzyloxycarbonylphenyl)glycine Methyl Ester

Prepared from Boc-R-4-trifluoromethanesulphonyloxy phenylglycine methyl ester and benzyl alcohol using Method

15 PAE-D.

^1H NMR

Intermediate PAE-13

20 Boc-R-(4-Carboxyphenyl)glycine Methyl Ester.

Boc-R-(4-benzyloxycarbonylphenyl)glycine methyl ester (500 mg) was dissolved in THF containing Pd/C 10% (100 mg) and hydrogenated at 1 atm for 2 h. Removal of the catalyst by filtration and evaporation of solvent gave Boc-R-(4-carboxy-phenyl)glycine methyl ester (330 mg, 87%).

^1H NMR

Intermediate PAE-14

30 Boc-R-(4-carboxamidophenyl)glycine Methyl Ester.

Method PAE-E

To a solution of Boc-R-(4-carboxyphenyl)glycine methyl ester (3.5 g) in DMF (30 mL) was added EDCI (2.60 g, 1.36 mmol) and HOBt (1.4 g, 10.4 mmol), and the mixture stirred for

10 min before cooling in a ice bath and bubbling in ammonia gas for 5 min. The mixture was stirred for 2 h at room temperature and then diluted with ethyl acetate and washed with water. The aqueous solution was extracted with a little ethyl acetate and the combined organics washed with brine. The organic solution was evaporated to an oil which was purified by flash chromatography (SiO_2 - dichloromethane/ethyl acetate 0 - 25%) to give Boc-R-(4-carboxamidophenyl)glycine methyl ester (1.7 g, 48%).

10

^1H NMR

Intermediate PAE-15

Boc-R-(4-methylcarboxamidophenyl)glycine Methyl Ester.

15

Prepared from Boc-R-(4-carboxyphenyl)glycine methyl ester and methylamine using Method PAE-E.

^1H NMR

20 Intermediate PAE-16

N-4-Methoxybenzoyl-N-2,4-dimethoxybenzyl-D,L-(quinolin-4-yl)glycine Methyl Ester.

Prepared from quinoline-4-carboxaldehyde using Method PAE-C.

25

^1H NMR

Intermediate PAE-17

Ethyl Boc-D,L-thiazol-2-ylglycine.

30

Prepared from ethyl hydroxyimino-thiazol-2-ylacetate using Method PAE-B. In this case, reaction with Zn/formic acid was conducted over 15 min.

¹NMR

IS-MS, m/e 287.0 (M+1)

Intermediate PAE-18

5 Ethyl Boc-D,L-isoquinolin-8-ylglycine.

Prepared from ethyl hydroxyimino-isoquinolin-8-ylacetate using Method PAE-B. In this case, reaction with Zn/formic acid was conducted over 30 min, followed by concentration and partitioning of the residue between 3/1 chloroform/isopropanol and satd aq. NaHCO₃. The Boc protection was carried out as previously described. Purification was performed using silica gel chromatography (Biotage Quad System) eluting with 10% ethyl acetate in methylene chloride.

15 ¹NMR

IS-MS, m/e 331.0 (M+1)

Analysis for C₁₈H₂₂N₂O₄:

Calcd: C, 65.44; H, 6.71; N, 8.48;

Found: C, 65.05; H, 6.67; N, 8.49.

20

Preparation of Intermediates PAA-1 - PAA-28

The following compounds were prepared according to the indicated method (Method PAA-A, Method PAA-B, Method PAA-C, Method PAA-D, Method PAA-E or Method PAA-F) from the indicated starting materials, unless otherwise described.

Intermediate PAA-1

Boc-D,L-(2-chlorophenyl)glycine.

Method PAA-A

30 2-Chlorobenzaldehyde (20 mmol, 2.252 mL) and 2,4-dimethoxybenzylamine (20 mmol, 3.004 mL) were added together and stirred for 2 hours. DCM (5 mL) was added and any water separated and removed. tert-Butyl isonitrile (20 mmol, 2.262 mL) was added and stirred for 10 min, followed by acetic acid

(20 mmol, 1.145 mL). Stirring was continued for 3 days. The reaction mixture was then treated with TFA (30 mL) and triethylsilane (5 mL). After 3 h the mixture was evaporated to dryness, 6 M HCl (100 mL) added, and the whole refluxed overnight at 130 °C, stirring rapidly. The mixture was allowed to cool and extracted with EtOAc (50 mL x 2); the aqueous fraction was evaporated to dryness and treated with 2 M NaOH solution. The mixture was extracted with EtOAc (50 mL x 2); excess boc anhydride (5.2 g) in dioxane (20 mL) was added to the aqueous fraction and stirred overnight. The mixture was extracted with diethyl ether (100 mL x 2), acidified to pH 1 (conc HCl) and extracted with EtOAc (50 mL x 2). The combined organic fractions were washed with water and evaporated to dryness under high vacuum. The product Boc -2-chlorophenylglycine (4.252 g, 74.5%)

^1H NMR ($\text{CD}_3\text{CN}/\text{D}_2\text{O}$) 7.3 (4H, m); 5.5 (1H, s); 1.3 (9H, s). MS 286 (M+1)

20 Intermediate PAA-1'

(R)-Benzyloxycarbonyl-(2-chlorophenyl)glycine.

Prepared from 2-chlorostyrene using the method of Sharpless et al J.A.C.S. (1998) Vol120 No.6 1207-1217.

25 Intermediate PAA-1, alternative preparation

Boc-D,L-(2-chlorophenyl)glycine.

Prepared from 2-chlorobenzaldehyde using method PAA-F. In this case, the reaction temperature was not controlled upon addition of 2-chlorobenzaldehyde and the reaction was allowed to stir for 2 h. Extraction of the intermediate aminonitrile was performed with ethyl ether in place of ethyl acetate and was further purified by addition of HCl gas to the ethereal extracts followed by decantation of the mother liquor to isolate the semisolid hydrochloride salt. BOC protection of

the amino acid was performed from 0 °C to room temperature over a period of one hour and the final extraction was performed with ethyl acetate in place of ethyl ether.

5 ¹H-NMR

IS-MS m/e 284 (M-1)

Intermediate PAA-2

Boc-D,L-(3-fluorophenyl)glycine.

10 Prepared from 3-fluorobenzaldehyde using Method PAA-A.

¹H NMR (CD₃CN/D₂O) 7.3 (1H, m), 7.1 (3H, m); 5.2 (1H, s); 1.3 (9H, s). MS 270 (M+1)

15 **Intermediate PAA-3**

Boc-D,L-(4-fluorophenyl)glycine.

Prepared from 4-fluorobenzaldehyde using Method PAA-A.

¹H NMR (CD₃CN/D₂O) 7.3 (2H, m); 6.9 (2H, m), 5.0 (1H, s); 1.3 (9H, s). MS 270 (M+1)

Intermediate PAA-4

Boc-D,L-(2-methylphenyl)glycine.

Prepared from 2-methylbenzaldehyde using Method PAA-A.

25

¹H NMR (CD₃CN/D₂O) 7.3 (4H, m); 5.5 (1H, s); 2.5 (3H, s); 1.3 (9H, s). MS 266 (M+1)

Intermediate PAA-5

30 **Boc-D,L-(3-thienyl)glycine.**

Prepared from 3-thiophenecarboxaldehyde using Method PAA-A.

¹H NMR (CD₃CN/D₂O) 7.5 (2H, m); 7.1 (1H, d); 5.3 (1H, s); 1.3 (9H, s). MS 258 (M+1)

Intermediate PAA-6

5 Boc-D,L-(2-fluorophenyl)glycine.

Was obtained by treating D,L-2-fluorophenylglycine (Aldrich) with Boc anhydride (1.1 eq) and 2 M NaOH (1 eq) in ethanol. Aqueous work up as described above yielded the protected amino acid.

10

¹H NMR

Intermediate PAA-7

Boc-D,L-(2-methoxyphenyl)glycine.

15 Prepared from 2-methoxybenzaldehyde using Method PAA-A.

¹H NMR

Intermediate PAA-7, alternative preparation

20 Boc-D,L-(2-methoxyphenyl)glycine.

Prepared from 2-methoxybenzaldehyde using method PAA-F. In this case, the reaction was cooled to 0 °C before addition of 2-methoxybenzaldehyde and was then allowed to stir at room temperature overnight. Extraction of the intermediate
25 aminonitrile was performed with ethyl ether in place of ethyl acetate and was further purified by addition of 1 M HCl in ethyl ether followed by filtration of the crystalline hydrochloride salt. BOC protection of the amino acid was performed from 0 °C to room temperature over a period of three
30 hours, and the final extraction was performed with dichloromethane in place of ethyl ether.

¹H-NMR

IS-MS m/e 280.1 (M-1)

Analysis for $C_{14}H_{19}NO_5$

Calcd: C, 59.78; H, 6.81; N, 4.98;

Found: C, 59.68; H, 6.78; N, 4.95.

5 Intermediate PAA-8

Boc-D,L-(2-trifluoromethyl)phenylglycine.

Prepared from 2-trifluoromethylbenzaldehyde using Method PAA-A.

10 1H NMR

Intermediate PAA-8, alternative preparation

Boc-D,L-(2-trifluoromethylphenyl)glycine.

Prepared from 2-trifluoromethylbenzaldehyde using method

- 15 PAA-F. In this case, the reaction temperature was not controlled upon addition of 2-trifluoromethylbenzaldehyde and the reaction was allowed to stir for 2 h. Extraction of the intermediate aminonitrile was performed with ethyl ether in place of ethyl acetate and was further purified by addition of
- 20 HCl gas to the ethereal extracts followed by decantation of the mother liquor to isolate the semisolid hydrochloride salt. BOC protection of the amino acid was performed from 0 °C to room temperature over a period of one hour and the final extraction was performed with ethyl acetate in place of ethyl
- 25 ether.

1H -NMR

IS-MS m/e 318 (M-1)

30 Intermediate PAA-9

Boc-D,L-(8-quinoliny)glycine.

Method PAA-B

To a stirring solution of Boc-D,L-(8-quinoliny)glycine

ethyl ester (2.29 g, 6.93 mmol) in 1,4-dioxane (11 mL) was added a solution of LiOH hydrate (0.32 g, 7.6 mmol) in water.

After 2 h, the solvents were removed in vacuo and the residue was dissolved in water and washed with diethyl ether. The aqueous phase was then acidified to pH 3 with solid citric acid and extracted with ethyl acetate. The organic phase was then washed with brine, dried with Na₂SO₄, filtered and concentrated to give 2.06 g (98%) of the title compound.

10 ¹H-NMR

IS-MS, m/e 303.0 (M+1)

Intermediate PAA-10

Boc-D,L-(5-quinolinyl)glycine.

15 Prepared from Boc-D,L-(5-quinolinyl)glycine ethyl ester using Method PAA-B.

¹H-NMR

IS-MS, m/e 303.0 (M+1)

20

Intermediate PAA-11

Boc-D-(3-bromophenyl)glycine.

Prepared from R-3-bromo-(1-t-butoxycarbonylamino-2-hydroxyethyl)benzene using Method PAA-C.

25

Melting Point = 130-132 °C with decomposition

¹H NMR (CDCl₃)

API-MS, m/e = 286 (M-CO₂H+1)

30 Intermediate PAA-12

Boc-D-(3-methoxycarbonylphenyl)glycine.

Method PAA-C

To a stirred solution of R-3-methoxycarbonyl-(1-t-butoxycarbonylamino-2-hydroxyethyl)benzene (338 mg, 1.14 mmol) in

acetone (7.2 mL) was added 5% NaHCO_3 (3 mL). The reaction mixture was cooled to 0 °C. To the stirred suspension was added KBr (14 mg, 0.12 mmol), TEMPO (181 mg, 1.16 mmol) and NaOCl dropwise (2.81 mL, 5.25%). After 1 h at 0 °C, TEMPO (136 mg, 0.88 mmol) and NaOCl (1.09 mL; 5.25%) were added. The reaction was stirred for a further 0.5 h at 0 °C and 5% NaHCO_3 (4.3 mL) was added. The reaction was allowed to warm to room temperature overnight. Acetone was removed under vacuum and the crude product was partitioned between ethyl acetate and water. The aqueous layer was washed with ethyl acetate (2x) and acidified to pH 5 with 10% citric acid and extracted with ethyl acetate (4x). The combined organic extracts were dried over MgSO_4 . Removal of solvent under vacuum gave the product (305 mg, 86%).

^1H NMR (CDCl_3)

API-MS, m/e = 254 ($\text{M}-\text{C}_4\text{H}_9+1$)

Intermediate PAA-13

Boc-D-(3-cyanophenyl)glycine.

Prepared from R-3-cyano-(1-t-butoxycarbonylamino-2-hydroxyethyl)benzene using Method PAA-C.

^1H NMR (CDCl_3)

API-MS, m/e = 221 ($\text{M}-\text{C}_4\text{H}_9+1$)

Intermediate PAA-14

Boc-D-(3-ethanesulfonylamino-phenyl)glycine.

To a stirring solution of 3-(ethanesulfonylamino-phenyl)glycine (20 g, 77.43 mmol) and sodium carbonate (8.2 g, 77.43 mmol) in 3:1 THF:water (200 mL) at 0 °C, was added di-tert-butyl dicarbonate (18.5 g, 85.17 mmol). After stirring for 30 min, the cold bath was removed; and after an additional 30 min at room temperature the solvent was removed; and the

residue was partitioned between ethyl acetate and water. The aqueous layer was acidified to pH 2 with KHSO_4 and extracted twice with ethyl acetate. The combined ethyl acetate extracts were washed with water, dried with Na_2SO_4 , filtered and concentrated in vacuo to give 17.51 g (63%) of a white solid.

^1H -NMR

IS-MS, m/e 357.0 (M-1)

10 Intermediate PAA-15

N-Boc-D,L-(5-thiazolyl)glycine.

To a r.b. flask (150 cm^3), was added Boc-D,L-(5-thiazolyl)glycine ethyl ester (7.00 g, 24.70 mmol) to ethanol (c.a. 100 cm^3) with stirring. 2 M Sodium hydroxide solution (25 cm^3 , 50 mmol) was added and allowed to stir for 1 h. Reaction monitored by TLC (60% hexane/ethyl acetate; s.m. r.f 0.5, prod. r.f. 0.). Reaction concentrated in vacuo and product taken up in saturated bicarbonate (c.a. 50 cm^3) and washed with ethyl acetate (c.a. 30 cm^3). Aqueous layer was acidified to pH 2 with concentrated hydrochloric acid and product extracted with 3:1 chloroform/isopropanol solution (c.a. 3x60 cm^3). The organic layer was dried over magnesium sulphate and evaporated to dryness to give an orange solid (4.47 g, 17.30 mmol) [74% Yield].

^1H NMR (CDCl_3); 1.35 (9H, s), 5.60 (1H, d), 5.83 (1H, d), 7.88 (1H, s), 8.80 (1H, s).

Intermediate PAA-16

30 N-Boc-D,L-(4-thiazolyl)glycine.

Method PAA-D

To a solution of N-Boc-D,L-(4-thiazolyl)glycine ethyl ester (0.700 g, 2.470 mmol) in methanol (c.a. 15 cm^3), was added 2 M sodium hydroxide (2.47 cm^3 , 4.940 mmol) and allowed

to stir for 90 min. The solution was concentrated *in vacuo* and taken up in water (c.a. 20 cm³). The aqueous solution was washed with ethyl acetate (c.a. 20 cm³), and then acidified to pH 2 with 5% hydrochloric acid solution (c.a. 50 cm³). The product was extracted with ethyl acetate (c.a. 3x30 cm³), dried over magnesium sulphate, and concentrated *in vacuo* to yield a pale yellow oil (0.582 g, 2.254 mmol) [91% yield].

¹H NMR (CDCl₃) 1.35 (9H, s), 5.5 (1H, d), 5.8 (1H, d), 7.35 (1H, d), 8.75 (1H, d), 9.8-10.2 (1H, br.).

Intermediate PAA-17

N-Boc-D,L-(2-methylthiazol-4-yl)glycine.

Prepared from N-Boc-D,L-(2-methylthiazol-4-yl)glycine ethyl ester using Method PAA-D.

¹H NMR (CDCl₃) 1.35 (9H, s), 2.6 (3H, s), 5.4 (1H, d), 5.9 (1H, s), 7.1 (1H, s).

20 Intermediate PAA-18

N-Boc-D,L-(2-Benzylloxycarbonylamino-4-thiazolyl)glycine.

Is prepared from D,L-(2-benzylloxycarbonylamino-4-thiazolyl)glycine. The benzylloxycarbonyl protecting group is removed from the thiazolyl amino group at a convenient point in the preparation of a final compound using a conventional method, such as, for example, heating a solution of an intermediate in HBr/acetic acid at 60 °C, followed by evaporation and a conventional isolation, such as by using SCX ion exchange chromatography.

30

D,L-(2-Benzylloxycarbonylamino-4-thiazolyl)glycine.

Was prepared by the method of Hardy, K.; Harrington, F. and Stachulski, A. - J. Chem. Soc. Perkin Trans I (1984) 1227-1235.

Intermediate PAA-19**Boc-R-(4-methoxycarbonylphenyl)glycine.**

To a solution of Boc-R-(4-methoxycarbonylphenyl)glycine
5 methyl ester (692 mg) in THF (10 mL) was added a solution of
lithium hydroxide hydrate (90 mg) in water (7 mL). The
mixture immediately became cloudy and over 15 min cleared.
After 30 min, tlc showed the reaction to be complete. Ethyl
acetate (20 mL) and water (20 mL) were added, and the aqueous
10 layer separated. The aqueous solution was acidified with 2 M
hydrochloric acid and extracted with ethyl acetate (3 x 20
mL). The organic solution was then washed with water x 2 and
brine x 2, dried with MgSO_4 and evaporated to give the mono-
ester (650 mg, 98%), pure by tlc.

15 ^1H NMR

Intermediate PAA-20**Boc-R-(4-Methoxyphenyl)glycine.**

20 Boc-R-(4-hydroxyphenyl)glycine methyl ester was converted
to Boc-R-4-methoxyphenylglycine using the alkylation method
described by Basak et al. (Tetrahedron Lett. 1998, 39 (27),
4883-4886), followed by hydrolysis of the methyl ester with
lithium hydroxide in aqueous THF.

25 ^1H NMR

Intermediate PAA-21

**N-4-Methoxybenzoyl-N-2,4-dimethoxybenzyl-D,L-(2-trifluoro-
30 methylphenyl)glycine.**

Prepared from N-4-methoxybenzoyl-N-2,4-dimethoxybenzyl-
D,L-(2-trifluoromethylphenyl)glycine methyl ester using Method
PAA-B (3 equivalents of LiOH hydrate).

¹H NMR

IS-MS, m/e 503.9 (m + 1)

Intermediate PAA-22

5 N-4-Methoxybenzoyl-N-2,4-dimethoxybenzyl-D,L-(thien-2-yl)-glycine.

Method PAA-E

To a solution of 2-thiopheneboronic acid (5.0 g, 39.0 mmol, 1 equiv) in 275 mL of methylene chloride at rt was added
10 3,4-dimethoxybenzylamine (5.89 mL, 39.0 mmol, 1 equiv) followed by glyoxylic acid monohydrate 3.6 g, 39 mmol, 1 equiv). The reaction was allowed to stir for 56 hours at rt after which time the resultant precipitate was filtered and washed with methylene chloride to afford 9.3 g (78%) of N-2,4-
15 dimethoxybenzyl-D,L-(thien-2-yl)glycine as an off-white solid (IS-MS, m/e 308 (m + 1)).

A portion of the solid (5.0 g, 16.3 mmol, 1 equiv.) was dissolved in acetone (20 mL) and 1 N sodium hydroxide (20 mL) at rt. To this solution was simultaneously added anisoyl
20 chloride (2.78 g, 16.3 mmol, 1 equiv.) in 20 mL of acetone and 2 N sodium hydroxide in dropwise fashion. After stirring at rt for 1 h, the reaction was cooled to 0 °C and was acidified to pH 2-3. Diethyl ether was added and the product was extracted into the organic phase. The combined organic phases
25 were washed with brine, dried over sodium sulfate, filtered, and concentrated to afford 5.1 g (71%) of the titled compound as a white solid.

IS-MS, m/e 440 (m + 1).

30

Intermediate PAA-23

N-Boc-N-2,4-dimethoxybenzyl-D,L-(thien-2-yl)glycine.

To a solution of N-2,4-dimethoxybenzyl-D,L-(thien-2-yl)glycine (1.0 g, 3.2 mmol, 1 equiv) in 6 mL of acetone and 6

mL of water at rt was added triethylamine (0.97 mL, 7.0 mmol, 2.1 equiv.) followed by addition of 2-(tert-butoxycarbonyloxyimino)-2-phenylacetonitrile (BOC-ON) (0.76 g, 3.1 mmol, 0.95 equiv). After stirring at rt overnight, the
5 reaction was diluted with water and washed with ether. The aqueous phase was then acidified with 0.5 M citric acid and the product was extracted into diethyl ether. The combined organic phases were washed with brine, dried over sodium sulfate, filtered, and concentrated to afford 0.38 g (29%) of
10 the titled compound as a crude yellow oil.

IS-MS, m/e 408 (m +1).

Intermediate PAA-24

15 Boc-D,L-isoquinolin-8-ylglycine.

Prepared from ethyl Boc-D,L-isoquinolin-8-ylglycine using Method PAA-B. The product was precipitated from a basic aqueous solution by adjusting the pH to 3 with solid citric acid.

20 ¹NMR

IS-MS, m/e 303.0 (M+1)

Analysis for C₁₆H₁₈N₂O₄·0.5 H₂O:

Calcd: C, 61.73; H, 6.15; N, 9.00;

Found: C, 61.62; H, 5.66; N, 8.84.

25

Intermediate PAA-25

Boc-D,L-Naphthalen-1-ylglycine.

Method PAA-F

Part A: D,L-Naphthalen-1-ylglycine hydrochloride.

30 To a solution of sodium cyanide (10.0 g, 0.22 mmol) in 40 mL of water was added ammonium chloride (11.4 g, 0.22 mmol), and the mixture was stirred until dissolution was complete. A solution of 1-naphthaldehyde (31.0 g, 0.22 mmol) in 40 mL of methanol was then added and the resultant mixture was allowed

to stir at room temperature for two days. An additional 150 mL of water was then added and the crude product was extracted into EtOAc. The combined organic layers were washed with water, dried over Na_2SO_4 , filtered and concentrated to afford 5 a crude oil. The crude residue was chromatographed over silica gel, eluting with with 10:1 EtOAc: CH_2Cl_2 , to give 35 g of a light brown oil. This material was then dissolved in 250 mL of 5 N HCl and was heated to reflux for 9 h. The reaction was allowed to cool to room temperature and the product was 10 allowed to crystallize overnight. Filtration of the mixture afforded 13.6 g (29%) of the title compound as light brown crystals.

^1NMR

15 IS-MS, m/e 201.9 (M+1)

Part B: Boc-D,L-Naphthalen-1-ylglycine.

To a solution of D,L-naphthalen-1-ylglycine hydrochloride (13.6 g, 57.2 mmol) and 2 N sodium hydroxide (57 mL, 115 mmol) 20 in 120 mL of 1,4-dioxane and 60 mL of water was added $(\text{Boc})_2\text{O}$ (15 g, 69 mmol). The reaction was allowed to stir at room temperature for 3 h after which time the solution was brought to pH 5 by addition of 1 N sulfuric acid. The product was then extracted into EtOAc; and the combined organic extracts 25 were dried over Na_2SO_4 , filtered, and concentrated to give 14 g (81%) of the title compound as a light brown foam.

^1NMR

IS-MS, m/e 300.1 (M-1)

30

Intermediate PAA-26

Boc-D,L-(2-methylthiophenyl)glycine.

To a solution of 2-(methylthio)benzaldehyde (15 g, 98.7 mmol) in 100 mL of ethanol was added ammonium carbonate (23.1

g, 296 mmol) and a solution of potassium cyanide (12 g, 148 mmol) in 100 mL water. The reaction was heated and stirred at 70 °C for 3 h after which time the reaction was concentrated under reduced pressure. The product was extracted into ethyl acetate; and the combined organic phases were washed with brine, dried over Na₂SO₄, filtered and concentrated. The resultant crude residue was taken up in 70 mL of ethyl acetate, and 70 mL of 5 N sodium hydroxide was added. The reaction was heated to reflux for three days after which time the ethyl acetate was removed under reduced pressure. To the aqueous mixture was sequentially added 100 mL of dioxane, Boc₂O (42 g, 192 mmol), and 100 mL of 2.5 N sodium hydroxide. The reaction was then heated at reflux for 48 h. After cooling to room temperature, the reaction was diluted with water and the aqueous phase was washed with ethyl ether. The aqueous layer was then acidified to pH 2 and the product was extracted into ethyl acetate. The combined organic extracts were washed with brine, dried over MgSO₄, filtered, and concentrated to afford 21.7 g of a crude residue.

Purification by silica gel chromatography (gradient elution, 97:2:1 to 95:4:1 dichloromethane:methanol:acetic acid) provided 5.0 g (17%) of the title compound.

¹H-NMR

ES-MS m/e 296 (M-1)

Intermediate PAA-27

Boc-D,L-(2-methylsulfonylphenyl)glycine.

To a solution of boc-D,L-(2-methylthiophenyl)glycine (4.5 g, 15.2 mmol) in 75 mL of methanol was added a solution of oxone (14 g, 23 mmol) in water. The reaction was stirred at room temperature for 2 h after which time the methanol was removed under reduced pressure. The product was extracted into ethyl acetate and the combined organic layers were washed

with brine, dried over MgSO_4 , filtered, and concentrated to afford 4.35 g (87%) of the title compound.

$^1\text{H-NMR}$

5 ES-MS m/e 230 ($\text{M}+1-\text{C}_5\text{H}_9\text{O}_2$)

Intermediate PAA-28

Boc-D,L-(benzo[b]thiophen-3-yl)glycine.

May be prepared by the method of Kukolja, S. et al.

10 *J. Med. Chem.* 1985, 28, 1886-1896.

General Experimental Procedures: Synthesis of Inhibitors

Coupling Method A:

15 The coupling of an amine and carboxylic acid to form an amide. A solution of the amine (1 equiv) and carboxylic acid (1.1 equiv) in a suitable solvent (DMF, and/or methylene chloride) is or was treated with diethyl cyanophosphonate (1.1 equiv) followed by addition of triethylamine or
20 diisopropylethyl amine (0 to 3 equiv) to the mixture. After completion of the reaction by thin-layer chromatography, the mixture was partitioned between a suitable solvent (EtOAc, and/or methylene chloride) and washed with 1 N NaOH, water, brine, and concentrated. The crude mixture was then purified,
25 as indicated, or used directly in subsequent transformations.

Coupling Method B:

The coupling of an amine and carboxylic acid to form an amide. A solution of the amine (1 equiv) and carboxylic acid
30 (1.1 equiv) in a suitable solvent (DMF, and/or methylene chloride) is or was treated with a carbodiimide-based dehydrating agent (e.g. DCC, or EDCI) (1.0 equiv). In general, addition of a benzotriazole-based reagent (e.g. HOBT or HOAT) (1 equiv) improved reaction yields. After completion of

the reaction by thin-layer chromatography, the mixture is or was partitioned between a suitable solvent (EtOAc, and/or methylene chloride) and washed with 1 N NaOH, water, brine, and concentrated. The crude mixture is or was then purified, 5 as indicated, or used directly in subsequent transformations.

Coupling Method C:

The coupling of an amine and acid chloride to form an amide. A solution of the amine (1 equiv) in an appropriate 10 solvent (chloroform, and/or methylene chloride) and pyridine (1-10 equiv) is or was treated with an acid chloride (1.1 equiv). After completion of the reaction by thin-layer chromatography, the mixture is or was partitioned between a suitable solvent (EtOAc, methylene chloride, and/or 15 chloroform) and washed with 1 N NaOH, water, brine, and concentrated. The crude mixture is or was then purified, as indicated, or used directly in subsequent transformations.

Deprotection Method A:

20 A mixture of 10% Palladium on carbon and the starting material in an appropriate solvent (EtOAc, EtOH, and/or HOAc) is or was placed under an atmosphere of hydrogen. Upon completion, the mixture is or was filtered and the resulting filtrate concentrated. The crude mixture is or was then 25 purified, as indicated, or used directly in subsequent transformations.

Alkylation Method A:

A solution of the starting material (1 equiv) in 5-10% 30 HOAc in methanol (anhydrous) is or was treated with the indicated aldehyde or ketone (2-10 equiv) followed by sodium cyanoborohydride (2-10 equiv). After completion, the mixture is or was concentrated and the residue was either partitioned between a suitable solvent (EtOAc, methylene chloride, and/or

chloroform) and washed with 1 N NaOH, water, brine, and concentrated or directly loaded onto an ion-exchange resin (SCX, Varian) and eluted with methanol followed by 2 N ammonia in methanol. Concentration of the later fractions afforded 5 the free base product. The crude mixture is or was then purified, as indicated, or used directly in subsequent transformations.

Preparation of Starting Materials

10

1-(Benzyloxycarbonyl-D-phenylglyciny)l)piperidine-4-methanol.

(Coupling Method A): A solution of benzyloxycarbonyl-D-phenylglycine (5.0 g, 17.5 mmol) and 4-piperidinemethanol (1.83 g, 15.9 mmol) in 90 mL of methylene chloride was cooled

15

in a methanol/ice bath and then treated with diethyl cyanophosphonate (2.67 mL, 17.5 mmol) followed by ethyl diisopropylamine (3.1 mL). After 5 h, the mixture was concentrated, diluted with EtOAc and saturated aqueous potassium carbonate, and the resulting layers separated. The 20 organic layer was washed (with aqueous potassium carbonate, 1 N HCl, brine), dried over magnesium sulfate, filtered, concentrated, and the residue purified by column chromatography (SiO₂: 70%-80% EtOAc:hexane), affording 1.54 g (25%) of the title compound.

25 ¹NMR

IS-MS, m/e 725 (M + 1).

Analysis for C₂₂H₂₆N₂O₄·0.15 H₂O:

Calcd: C, 68.6; H, 6.9; N, 7.3;

Found: C, 68.5; H, 6.9; N, 7.1.

30

1-(D-Phenylglyciny)l)piperidine-4-methanol.

Using Deprotection Method A, 1-(benzyloxycarbonyl-D-phenylglyciny)l)piperidine-4-methanol (3.93 g, 29.5 mmol) and 10% palladium on carbon (1.30 g) in 2:1 EtOAc:EtOH (75 mL)

afforded 2.31 g (88%) of the title compound.

¹NMR

IS-MS, m/e 249 (M+1).

5 Preparation of Intermediates A-1 - A-3

Intermediate A-1

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-methanol.

Using Coupling Method C, 1-(D-phenylglyciny)l)piperidine-4-methanol (1.23 g, 4.96 mmol) and p-anisoyl chloride (0.888 g, 5.21 mmol) afforded, after purification by column chromatography (SiO₂: 1:1 to 1:9 hexanes:EtOAc), 1.26 g (66%) of the title compound.

¹NMR

15 IS-MS, m/e 383 (M+1).

Intermediate A-2

1-(Indole-6-carbonyl-D-phenylglyciny)l)piperidine-4-methanol.

(Coupling Method B): A solution of 1-(D-phenylglyciny)l)

20 piperidine-4-methanol (500 mg, 2.02 mmol), indole-6-carboxylic acid (325 mg, 2.02 mmol), and 1-hydroxy-7-azabenzotriazole (275 mg, 2.02 mmol) in 10 mL of DMF was treated with DCC (415 mg, 2.02 mmol). After 15 h, the mixture was concentrated and the residue dissolved in EtOAc. The organic layer was washed
25 (with 2 N NaOH, water, brine), dried over magnesium sulfate, and concentrated. The residue was purified by column chromatography (SiO₂: EtOAc), affording 780 mg (98%) of the title compound.

¹NMR

30 IS-MS, m/e 383 (M + 1).

Intermediate A-3

1-(3-Chloroindole-6-carbonyl-D-phenylglyciny)l)piperidine-4-methanol.

Using Coupling Method B, 1-(D-phenylglyciny)l)piperidine-4-methanol (9.90 g, 39.9 mmol) and 3-chloroindole-6-carboxylic acid (8.56 g, 43.9 mmol) afforded, after purification by column chromatography (SiO₂: 95% EtOAc in hexane), 8.95 g (53%) of the title compound.

¹NMR

IS-MS, m/e 426 (M + 1).

Preparation of Intermediate B-1

Intermediate B-1

1-(4-Methoxybenzoyl-D-phenylglyciny)l)piperidine-4-carboxaldehyde

A solution of 1-(4-methoxybenzoyl-D-phenylglyciny)l)-piperidine-4-methanol (0.800 g, 2.08 mmol) and N-methylmorpholine oxide (0.366 g, 3.13 mmol) in methylene chloride (15 mL) was treated with tetrapropylammonium perruthenate (TPAP, 2 mg). After 14 h, the mixture was treated with additional TPAP (5 mg). After 20 h, the mixture was treated with additional TPAP (5 mg). After 32 h, the mixture was loaded directly onto a column and purified by column chromatography (SiO₂: 1:1 to 1:4 hexanes:EtOAc) affording 0.286 g (36%) of the title compound.

¹NMR

IS-MS, m/e 381 (M+1).

Preparation of Intermediates C-2 - C-4

Intermediate C-2

1-(Indole-6-carbonyl-D-phenylglyciny)l)-4-(methylsulfonyloxymethyl)piperidine.

A solution of 1-(indole-6-carbonyl-D-phenylglyciny)l)-piperidine-4-methanol (2.82 g, 7.21 mmol) and triethylamine (2.0 mL) in 40 mL of methylene chloride was treated with

methanesulfonyl chloride (1.1 mL, 14.0 mmol). After 2 h, mixture was concentrated and the residue purified by column chromatography (SiO₂: 90% EtOAc in hexane) to afford 2.80 g (83%) of the title compound.

5 ¹NMR

IS-MS, m/e 470 (M + 1).

Intermediate C-3

1- (3-Chloroindole-6-carbonyl-D-phenylglyciny)l)-4-

10 (methylsulfonyloxymethyl)piperidine.

Using a procedure similar to that described above for Intermediate C-2, 1-(3-chloroindole-6-carbonyl-D-phenylglyciny)l)piperidine-4-methanol (8.95 g, 21.0 mmol) afforded, after purification by column chromatography (SiO₂: 80% to 99% EtOAc in hexane), 2.58 g (24%) of the title compound.

15 ¹NMR

IS-MS, m/e 504 (M + 1).

(Protected) Intermediate C-4

20 1-(1-Methylsulfonyl-3-chloroindole-6-carbonyl-D-phenylglyciny)l)-4-(methylsulfonyloxymethyl)piperidine.

Using a procedure similar to that described above for Intermediate C-3, 1-(3-chloroindole-6-carbonyl-D-phenylglyciny)l)piperidine-4-methanol (8.95 g, 21.0 mmol) afforded, as a minor side product after purification by column chromatography (SiO₂: 80% to 99% EtOAc in hexane), 940 mg (8%) of the title compound.

25 ¹NMR

IS-MS, m/e 582 (M+1)

30

Preparation of Intermediate D-1

4-(Piperidin-1-ylmethyl)piperidine bis-Hydrochloride Salt.

A solution of 4-(piperidin-1-ylmethyl)pyridine (10.0 g,

0.056 mol; prepared using a similar procedure to that described in US430491) and platinum oxide (1.5 g) in 55 mL of ethanol and 18 mL of 12 N HCl was placed under an atmosphere of hydrogen (4.1 bar, 60 psig). After 15 h, the mixture was filtered, concentrated, and residue triturated with methanol affording 9.0 g of the title compound as a bis hydrochloride salt.

¹NMR

IS-MS, m/e 183 (M + 1).

10

1-(Benzyloxycarbonyl-D-phenylglyciny)-4-[(piperidin-1-yl)-methyl]piperidine.

Using Coupling Method A, benzyloxycarbonyl-D-phenylglycine (17.1 g, 60.1 mmol) and 4-(piperidin-1-ylmethyl)-piperidine (10.0 g, 54.6 mmol) afforded, after purification by column chromatography (SiO₂: 2% to 3% [2 N ammonia in methanol]:methylene chloride) 10.7 g (43%) of the title compound.

¹NMR

20 IS-MS, m/e 451 (M + 1).

Analysis for C₂₇H₃₅N₃O₃:

Calcd: C, 72.1; H, 7.8; N, 9.3;

Found: C, 72.0; H, 7.7; N, 9.4.

25 Intermediate D-1.

1-(D-Phenylglyciny)-4-[(piperidin-1-yl)methyl]piperidine.

Using Deprotection Method A, 1-(benzyloxycarbonyl-D-phenylglyciny)-4-[(piperidin-1-yl)methyl]piperidine (10.7 g, 23.8 mmol) and 10% palladium on carbon (1.35 g) in 200 mL of 1:1 EtOH:EtOAc afforded the title compound.

¹NMR

IS-MS, m/e 315 (M + 1).

Preparation of Intermediate D-2

1- [Boc-D,L- (2-methoxyphenyl)glyciny] -4- [(piperidin-1-yl) -
methyl]piperidine.

Using Coupling Method B, Boc-D,L-(2-methoxyphenyl)glycine
5 (2.0 g, 7.11 mmol) and 4-(piperidin-1-ylmethyl)piperidine
(1.43 g, 7.82 mmol) with triethylamine (1 mL, 7.8 mmol)
afforded, after purification by column chromatography (SiO₂;
0-2% [2 N ammonia in methanol]:methylene chloride), 1.59 g
(50%) of the title compound.

10 ¹NMR

IS-MS, m/e 446.1 (M+1)

Intermediate D-2.

1- [D,L- (2-Methoxyphenyl)glyciny] -4- (piperidin-1-ylmethyl) -
15 piperidine.

To a stirring solution of 1-(Boc-D,L-2-methoxyphenyl) -
glyciny] -4- (piperidin-1-ylmethyl)piperidine (1.52 g, 3.42
mmol) in methylene chloride (15 mL) was added anisole (2.6 mL,
24 mmol) followed by trifluoroacetic acid (5.3 mL). After
20 stirring for 3 h, the solvent was removed in vacuo and the
residue was dissolved a minimal volume of methanol and loaded
onto an SCX column. The column was eluted with methanol,
followed by 2 N ammonia in methanol. The product containing
fractions were combined and concentrated in vacuo to give 1.2
25 g (100%) of the title compound.

¹NMR

IS-MS, m/e 346.1 (M+1)

Preparation of Intermediate D-3

30

1- (Boc-D,L-thiazol-5-ylglyciny] -4- [(piperidin-1-yl)methyl] -
piperidine.

To a solution of N-Boc-D,L-(5-thiazolyl)glycine (1.36 g,
5.25 mmol), HOAT (787 mg 5.78mmol), 4-(1-piperi-

dinylmethyl)piperidine (1.35 g 5.25 mmol) and triethylamine (1.61 mL, 11.6 mmol) in DMF (42 mL) was added EDCI (1.11 g, 5.78 mmol) and the mixture stirred at room temperature for 20 h. The solvent was removed *in vacuo*, the residues taken up in chloroform:isopropyl alcohol (2:1) and washed with water and satd sodium bicarbonate. The aqueous phase was back extracted with chloroform:isopropyl alcohol (2:1) (x2), and the combined organic extracts were dried (MgSO_4) and concentrated *in vacuo* to an orange-brown oil. The crude reaction product thus obtained was carried on to the next step without further purification.

Intermediate D-3.

1- (D,L-thiazol-5-ylglyciny)l)-4-[(piperidin-1-yl)methyl]-piperidine.

To a stirred solution of crude 1-(Boc-D,L-thiazol-5-ylglyciny)l)-4-[(piperidin-1-ylmethyl)piperidine] (circa 5.25 mmol) and anisole (11.4 mL) in dichloromethane (42 mL) at room temperature was added TFA (10.5 mL) and the mixture stirred at room temperature for 20 h before concentrating *in vacuo*. The product was isolated using SCX ion exchange chromatography.

¹H NMR

Preparation of Intermediate D-4

1- [N-Boc-D,L-(2-benzyloxycarbonylaminothiazol-4-yl)glyciny]l)-4-[(piperidin-1-yl)methylpiperidine].

This was prepared by coupling N-Boc-D,L-(2-benzyloxy-carbonylamino-4-thiazolyl)glycine and 4-(1-piperidinyl-methyl)piperidine as described above in the corresponding step in the preparation of Intermediate D-3.

Intermediate D-4.

1- (D,L-2-Aminothiazol-4-ylglyciny)l)-4-[(piperidin-1-yl)-

methyl]piperidine].

A stirred solution of 1-[N-Boc-D,L-(2-benzyloxy-carbonylamino-4-thiazolyglycyl)-4-[(piperidin-1-yl)-methyl]piperidine] (crude, circa 5.25 mmol) in a mixture of
5 HBr-acetic acid (50%, 35 mL) and acetic acid (70 mL) was heated at 60 °C for 7 h, cooled and then concentrated in vacuo.

The product was isolated using SCX ion exchange chromatography.

¹NMR

10

Preparation of Intermediate D-5

1-(Boc-D,L-2-methylthiazol-4-ylglycyl)-4-[(1-piperidinyl)-methyl]piperidine.

15

This was prepared by coupling N-Boc-D,L-(2-methylthiazol-4-ylglycine and 4-(1-piperidinylmethyl)piperidine as described above in the corresponding step in the preparation of Intermediate D-3.

20 Intermediate D-5.

1-(D,L-2-Methylthiazol-4-ylglycyl)-4-[(piperidin-1-yl)-methyl]piperidine.

This was prepared by deprotection of 1-(Boc-D,L-2-methylthiazol-4-ylglycyl)-4-[(piperidin-1-yl)methyl]-

25 piperidine as above in the corresponding step in the preparation of Intermediate D-3.

¹NMR

Preparation of Intermediate D-6

30

1-Boc-4-[(2,3,5,6-tetrachloropyrid-4-ylamino)methyl]piperidine.

A mixture of 1-Boc-4-(aminomethyl)piperidine (2.0 g, 9.33 mmol), pentachloropyridine (2.58 g, 10.3 mmol) and

potassium carbonate (2.58 g, 18.7 mmol) was heated at 60 °C in DMF (22mL) for 2 h. Excess DMF was removed *in vacuo* and the residue taken in to water/ethyl acetate. The aqueous solution was extracted with ethyl acetate (x3) and the combined organic solution dried (MgSO₄) and concentrated *in vacuo*. Flash chromatography (SiO₂, hexane/diethyl ether 30% to 60% gave the product as a white solid (2.5g, 62%).

¹NMR

10 1-Boc-4-[(pyrid-4-ylamino)methyl]piperidine.

To a stirred solution of 1-Boc-4-[(2,3,5,6-tetrachloro-pyrid-4-ylamino)methyl]piperidine (2.5 g, 5.83 mmol) in a mixture of methanol (23 mL) and THF (23 mL) was added sodium methoxide (1.89 g, 35 mmol) and 5% palladium on carbon

15 (1.24 g). The mixture was hydrogenated for 30 h and the catalyst was filtered off and the solution concentrated *in vacuo*. The residue was taken up in ethyl acetate and washed with water and satd sodium bicarbonate. The aqueous washes were back extracted with ethyl acetate, and the combined
20 organics dried (MgSO₄) and concentrated *in vacuo* to give the product (1.0 g, 59%).

¹NMR

4-[(Pyrid-4-ylamino)methyl]piperidine Dihydrochloride.

25 To a stirred solution of 1-Boc-4-[(pyrid-4-ylamino)-methyl]piperidine (1.0 g, 3.43 mmol) in methanol (100 mL) was added HCl (1 M in ether, 34 mL), and the mixture stirred overnight. The solvent was removed *in vacuo*, and the residue taken up in water and freeze dried to give the product as a
30 yellow solid (800 mg, 88%).

1-(Boc-D-phenylglycyl)-4-[(pyrid-4-ylamino)methyl]piperidine Trifluoroacetic Acid Salt.

To a mixture of Boc-D-phenylglycine (276 mg, 1.1 mmol),

4-[(pyrid-4-ylamino)methyl]piperidine dihydrochloride (290 mg, 1.1 mmol), HOAT (165 mg, 1.21 mmol) and triethylamine (367 mg, 3.63 mmol) in DMF (9 mL) was added EDCI (232 mg, 1.21 mmol), and the mixture stirred for 7 h at ambient temperature and 64 h at 5 °C. The solvent was removed *in vacuo* and the residue partitioned between dilute aqueous sodium bicarbonate and 2:1 chloroform:isopropanol. The aqueous solution was back extracted with 2:1 chloroform:isopropanol three times and the combined extracts dried (MgSO₄). The solvent was removed *in vacuo*, and the crude product purified by RPHPLC to give product (500 mg) as a TFA salt.

Intermediate D-6.

1-(D-Phenylglycynyl)-4-[(pyrid-4-ylamino)methyl]piperidine.

To 1-(Boc-D-phenylglycynyl)-4-[(pyrid-4-ylamino)-methyl]piperidine TFA salt (500 mL, 1.1 mmol) dissolved in DCM (9 mL) and anisole (1.79 mL, 16.5 mmol) was added TFA (2.3 mL), and the mixture stirred for 3 h. The solvent was removed *in vacuo*, and the residue purified by SCX ion exchange chromatography to give the product (290 mg, 81% over two steps).

Preparation of Examples 1-5

General Procedure: Unless otherwise indicated, the product of Examples 1-5 was obtained from the indicated amine and 1-(4-methoxybenzoyl-D-phenylglycynyl)piperidine-4-carboxaldehyde (Intermediate B-1) using Alkylation Method A.

Example 1.

1-[(4-Methoxybenzoyl-D-phenylglycynyl)]-4-[(isopropylamino)-methyl]piperidine Hydrochloride.

1-(4-Methoxybenzoyl-D-phenylglycynyl)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and isopropylamine

afforded, after treatment of the isolated product with excess hydrochloric acid in methanol and concentration, 37 mg of the title compound as a hydrochloride salt.

¹NMR

5 IS-MS, m/e 424 (M+1)

Example 2.

1-(4-Methoxybenzoyl-D-phenylglyciny1)-4-[(dimethylamino)-methyl]piperidine.

10 1-(4-Methoxybenzoyl-D-phenylglyciny1)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and dimethylamine afforded 25 mg (47%) of the title compound.

¹NMR

IS-MS, m/e 410 (M+1)

15

Example 3.

1-[(4-Methoxybenzoyl-D-phenylglyciny1)]-4-[(N,N-diethyl-amino)methyl]piperidine Hydrochloride.

1-(4-Methoxybenzoyl-D-phenylglyciny1)piperidine-4-
20 carboxaldehyde (0.050 g, 0.131 mmol) and diethylamine afforded, after treatment of isolated product with excess hydrochloric acid in methanol and concentration, 42 mg of the title compound as a hydrochloride salt.

¹NMR

25 IS-MS, m/e 438 (M+1)

Example 4.

1-[(4-Methoxybenzoyl-D-phenylglyciny1)]-4-[(1-pyrrolidinyl)-methyl]piperidine.

30 1-(4-Methoxybenzoyl-D-phenylglyciny1)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and pyrrolidine afforded 27 mg (47%) of the title compound.

¹NMR

IS-MS, m/e 436 (M+1)

Example 5.

1-[(4-Methoxybenzoyl-D-phenylglyciny)]-4-[(3-pyrrolin-1-yl)-methyl]piperidine Hydrochloride.

- 5 1-(4-Methoxybenzoyl-D-phenylglyciny)piperidine-4-carboxaldehyde (0.050 g, 0.131 mmol) and 3-pyrroline afforded, after treatment of isolated product with excess hydrochloric acid in methanol and concentration, 43 mg of the title compound as a hydrochloride salt.

10 ¹NMR

IS-MS, m/e 434 (M+1)

Preparation of Examples 6-18

- 15 **General Procedure:** Unless otherwise indicated, the product of Examples 6-18 was obtained from the indicated amine and 1-(indole-6-carbonyl-D-phenylglyciny)-4-(methylsulfonyloxymethyl)piperidine (Intermediate C-2) or 1-(3-chloroindole-6-carbonyl-D-phenylglyciny)-4-(methylsulfonyloxy-
- 20 methyl)piperidine (Intermediate C-3) using a procedure similar to that described in Example 6.

Where indicated, free bases were obtained by using SCX ion exchange chromatography; hydrochloride salts were obtained by dissolving the free base in ethyl acetate, treating with

25 1 M HCl in ether, removing the solvent *in vacuo*, redissolving the resulting HCl salt in water/acetonitrile and lyophilising.

Example 6.

- 1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(isopropylamino)-
- 30 methyl]piperidine Trifluoroacetate Salt.

A solution of 1-(indole-6-carbonyl-D-phenylglyciny)-4-(methylsulfonyloxymethyl)piperidine (100 mg, 0.213 mmol) and isopropylamine (0.18 mL, 2.1 mmol) in 1 mL of THF was treated with potassium carbonate (60 mg) and sodium iodide (32 mg,

0.21 mmol), and the mixture was heated at reflux. After 16 h, the mixture was concentrated and the residue purified by rpHPLC chromatography affording 24 mg (21%) of the title compound as a trifluoroacetate salt.

5 ¹NMR

IS-MS, m/e 433 (M+1).

Example 6a.

1- (Indole-6-carbonyl-D-phenylglyciny)l-4-[(isopropylamino)-
10 methyl]piperidine Hydrochloride Salt.

The trifluoroacetate salt was converted to the hydrochloride salt via the free base as described above.

ES-MS, m/e 433 (M+1).

15 **Example 7.**

1- (Indole-6-carbonyl-D-phenylglyciny)l-4-[4-(pyrrolidin-1-yl)methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (200 mg, 0.426 mmol) and pyrrolidine (0.85 mmol) afforded, after purification by rpHPLC

20 chromatography, 38 mg (16%) of the title compound as a trifluoroacetate salt.

¹NMR

Example 7a.

25 1- (Indole-6-carbonyl-D-phenylglyciny)l-4-[4-(pyrrolidin-1-yl)-methyl]piperidine Hydrochloride Salt.

The trifluoroacetate salt was converted to the hydrochloride salt via the free base as described above.

ES-MS, m/e 445 (M+1).

30

Example 8.

1- (Indole-6-carbonyl-D-phenylglyciny)l-4-[(diethylamino)-methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and diethylamine

(2.1 mmol) afforded, after purification by rpHPLC chromatography, 59 mg (49%) of the title compound as a trifluoroacetate salt.

¹NMR

5 IS-MS, m/e 447 (M+1).

Example 8a.

1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(diethylamino)-methyl]piperidine Hydrochloride salt.

10 The trifluoroacetate salt was converted to the hydrochloride salt via the free base as described above.
ES-MS, m/e 447 (M+1).

Example 9.

15 1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(piperidin-1-yl)-methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and piperidine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 16 mg (13%) of the title compound as a trifluoroacetate salt.

20 ¹NMR

IS-MS, m/e 459 (M+1).

Example 9a.

25 1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(piperidin-1-yl)-methyl]piperidine Hydrochloride Salt.

The trifluoroacetate salt was converted to the hydrochloride salt via the free base as described above.

ES-MS, m/e 449 (M+1).

30 Example 10.

1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(4-hydroxypiperidin-1-yl)methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and 4-hydroxypiperidine (2.3 mmol) afforded, after purification by rpHPLC

chromatography, 32 mg (25%) of the title compound as a trifluoroacetate salt.

¹NMR

IS-MS, m/e 475 (M+1).

5

Example 10a.

1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(4-hydroxypiperidin-1-yl)methyl]piperidine Hydrochloride Salt.

The trifluoroacetate salt was converted to the

10 hydrochloride salt via the free base as described above.

ES-MS, m/e 475 (M+1).

Example 11.

15 **1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(piperazin-1-yl)-methyl]piperidine Trifluoroacetate Salt.**

Intermediate C-2 (100 mg, 0.213 mmol) and piperazine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 85 mg (70%) of the title compound as a trifluoroacetate salt.

¹NMR

20 IS-MS, m/e 460 (M+1).

Example 12.

1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(4-methylpiperazin-1-yl)methyl]piperidine Trifluoroacetate Salt.

25 Intermediate C-2 (100 mg, 0.213 mmol) and 1-methylpiperazine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 66 mg (53%) of the title compound as a trifluoroacetate salt.

¹NMR

30 IS-MS, m/e 474 (M+1).

Example 13.

1- (Indole-6-carbonyl-D-phenylglyciny)l)-4- [(3-hydroxy-pyrrolidin-1-yl)methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and 3-hydroxy-
5 pyrrolidine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 36 mg (30%) of the title compound as a trifluoroacetate salt.

¹NMR

IS-MS, m/e 461 (M+1).

10

Example 14.

1- (Indole-6-carbonyl-D-phenylglyciny)l)-4- [(R) - (3-hydroxy-methylpyrrolidin-1-yl)methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and (R)-3-
15 hydroxymethylpyrrolidine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 32 mg (26%) of the title compound as a trifluoroacetate salt.

¹NMR

IS-MS, m/e 475 (M+1).

20

Example 14a.

1- (Indole-6-carbonyl-D-phenylglyciny)l)-4- [(R) - (3-hydroxy-methylpyrrolidin-1-yl)methyl]piperidine Hydrochloride Salt.

The trifluoroacetate salt was converted to the
25 hydrochloride salt via the free base as described above.

ES-MS, m/e 475 (M+1).

Example 15.

1- (Indole-6-carbonyl-D-phenylglyciny)l)-4- [(S) - (3-hydroxy-methylpyrrolidin-1-yl)methyl]piperidine Trifluoroacetate Salt.
30

Intermediate C-2 (100 mg, 0.213 mmol) and (S)-3-hydroxymethylpyrrolidine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 36 mg (29%) of the title compound as a trifluoroacetate salt.

¹NMR

IS-MS, m/e 475 (M+1).

Example 16.

- 5 1-(Indole-6-carbonyl-D-phenylglyciny)-4-[(4-hydroxymethyl-piperidin-1-yl)methyl]piperidine Trifluoroacetate Salt.

Intermediate C-2 (100 mg, 0.213 mmol) and 4-hydroxy-methylpiperidine (2.3 mmol) afforded, after purification by rpHPLC chromatography, 28 mg (22%) of the title compound as a
10 trifluoroacetate salt.

¹NMR

IS-MS, m/e 489 (M+1).

Example 17.

- 15 1-(3-Chloroindole-6-carbonyl-D-phenylglyciny)-4-[(piperidin-1-yl)methyl]piperidine Hydrochloride Salt.

Intermediate C-3 (1.0 g, 2.0 mmol) and piperidine (6.0 mmol) afforded, after purification by column chromatography (SiO₂: 2% to 4% [2 N ammonia in
20 methanol]:methylene chloride) and salt formation with hydrochloric acid, 443 mg (41%) of the title compound as a hydrochloride salt.

¹NMR

IS-MS, m/e 493 (M+1).

25

Example 18.

1-(3-Chloroindole-6-carbonyl-D-phenylglyciny)-4-[(3-hydroxypiperidin-1-yl)methyl]piperidine.

Intermediate C-3 (1.50 g, 3.0 mmol) and 3-hydroxy-
30 piperidine (9.0 mmol) afforded, after purification by column chromatography (SiO₂: 2% to 3% [2 N ammonia in methanol]:methylene chloride), 1.10 g of the title compound.

¹NMR

IS-MS, m/e 507 (M+1).

Preparation of Examples 19-21

General Procedure: Unless otherwise indicated, the product of
5 Examples 19-21 was obtained from the indicated carboxylic acid
and 1-(D-phenylglyciny1)-4-[(piperidin-1-yl)methyl]piperidine
(Intermediate D-1) using a procedure similar to that described
in Example 19.

10 Example 19.

1-(Indole-6-carbonyl-D-phenylglyciny1)-4-[(piperidin-1-yl)-
methyl]piperidine.

(Coupling Method B): A solution of 1-(D-phenylglyciny1)-4-
[(piperidin-1-yl)methyl]piperidine (4.00 g, 12.7 mmol) and
15 indole-6-carboxylic acid (2.25 g, 13.9 mmol) in 60 mL of DMF
was treated with HOBT (1.88 g, 13.9 mmol), the mixture cooled
to 0 °C, and then treated with DCC (2.87 g, 13.9 mmol). After
15 h, the mixture was diluted with EtOAc, filtered, and the
filtrate concentrated. The residue was purified by column
20 chromatography (SiO₂; 2% to 3.5% [2 N ammonia in
methanol]:methylene chloride) to afford 3.1 g (53%) of the
title compound.

¹NMR

IS-MS, m/e 459 (M+1).

25

Example 20.

1-(3-Methylindole-6-carbonyl-D-phenylglyciny1)-4-[(piperidin-
1-yl)methyl]piperidine Hydrochloride Salt.

Intermediate D-1 (2.00 g, 6.35 mmol) and 3-methylindole-
30 6-carboxylic acid (1.22 g, 6.98 mmol) afforded, after
formation of the hydrochloride salt, 1.29 g (41%) of the title
compound.

¹NMR

IS-MS, m/e 473 (M+1).

Analysis for $C_{29}H_{36}N_4O_2 \cdot HCl \cdot 0.75 H_2O$:

Calcd: C, 66.7; H, 7.4; N, 10.7;

Found: C, 66.7; H, 7.2; N, 10.8.

5 **Example 21.**

1- (5-Chloroindole-2-carbonyl-D-phenylglyciny1)-4-[(piperidin-1-yl)methyl]piperidine Hydrochloride Salt.

Intermediate D-1 and 5-chloro-1H-indole-2-carboxylic acid (1.36 g, 6.98 mmol) afforded, after formation of the hydrochloride salt, 1.52 g (45%) of the title compound.

¹NMR

IS-MS, m/e 493 (M+1).

Analysis for $C_{28}H_{33}ClN_4O_2 \cdot HCl \cdot 0.75 H_2O$:

Calcd: C, 61.9; H, 6.6; N, 10.3;

15 Found: C, 61.8; H, 6.4; N, 10.4.

Example 22.

1- [4-Methoxybenzoyl-D,L-(2-methoxyphenyl)glyciny1]-4-[(piperidin-1-yl)methyl]piperidine hydrochloride.

20 Using Coupling Method B, 4-methoxybenzoic acid (0.29 g, 1.89 mmol) and 1- [D,L-(2-methoxyphenyl)glyciny1]-4-[(piperidin-1-yl)methyl]piperidine (0.59 g, 1.7 mmol) afforded, after purification by column chromatography (SiO₂; 0-4% [2 N ammonia in methanol]:methylene chloride), 0.47 g of 25 1- [4-methoxybenzoyl-D,L-(2-methoxyphenyl)glyciny1]-4-[(piperidin-1-yl)methyl]piperidine. The free base was dissolved in methylene chloride, and 1 M HCl in diethyl ether (1 mL, 1 mmol) was added with stirring. The mixture was partially concentrated and filtered to give 0.34 g (39%) of 30 the title compound followed by a second crop of 0.12 g (14%).

¹NMR

IS-MS, m/e 480.1 (M+1)

Analysis for $C_{28}H_{37}N_3O_4 \cdot 0.95 HCl \cdot 1.0 H_2O$

Calcd: C, 63.18; H, 7.57; N, 7.90; Cl, 6.33;

Found: C, 63.43; H, 7.41; N, 7.68; Cl, 5.93.

5 Example 23.

1-[Indole-6-carbonyl-D,L-(2-methoxyphenyl)glycinyll]-4-
[(piperidin-1-yl)methyl]piperidine hydrochloride.

Using Coupling Method B, indole-6-carboxylic acid
(0.29 g, 1.89 mmol) and 1-[D,L-(2-methoxyphenyl)glycinyll]-4-
10 [(piperidin-1-yl)methyl]piperidine (0.59 g, 1.7 mmol)
afforded, after purification by column chromatography (SiO_2 ;
0-4% [2 N ammonia in methanol]:methylene chloride), 0.36 g of
1-[indole-6-carbonyl-D,L-(2-methoxyphenyl)glycinyll]-4-
[(piperidin-1-yl)methyl]piperidine. The free base was
15 dissolved in methylene chloride, and 1 M HCl in diethyl ether
(1 mL, 1 mmol) was added with stirring. The mixture was
partially concentrated and filtered to give 0.21 g (24%) of
the title compound followed by a second crop of 0.065 g (7%).
1NMR

20 IS-MS, m/e 489.0 (M+1)

Analysis for $C_{29}H_{36}N_4O_3 \cdot 1.0HCl \cdot 1.5H_2O$

Calcd: C, 63.09; H, 7.30; N, 10.15; Cl, 6.42;

Found: C, 62.99; H, 7.06; N, 9.90; Cl, 6.27.

25 Example 24.

1-(4-Methoxybenzoyl-D,L-thiazol-5-ylglycinyll)-4-[(piperidin-1-yl)
methyl]piperidine Hydrochloride Salt.

To a stirred solution of 4-methoxybenzoic acid (797 mg,
5.24 mmol), 1-(D,L-thiazol-5-ylglycinyll)-4-[(piperidin-1-yl)-
30 methyl]piperidine (circa 5.24 mmol) and HOAT (786 mg, 5.76
mmol) in DMF (40 mL) was added EDCI (1.11 g, 5.76 mmol). The
mixture was stirred at room temperature for 20 h and the
solvent removed in vacuo. The residues taken up in
chloroform: isopropyl alcohol (2:1) and washed with satd

sodium bicarbonate. The aqueous phase was back extracted with chloroform: isopropyl alcohol (2:1) (x3), and the combined organic extracts were dried (MgSO_4) and concentrated in vacuo.

Half of the crude product was purified by preparative RPHPLC; and the product fractions concentrated, taken up in chloroform: isopropyl alcohol (2:1), washed with satd sodium bicarbonate, dried (MgSO_4) and concentrated in vacuo. The free base thus obtained was dissolved in methanol and treated with 2 equivalents of HCl in ether and evaporated to dryness.

10 The residue was dissolved in water/acetonitrile and freeze dried. Yield 672 mg.

^1NMR

LCMS, m/e 457 (M+1)

15 **Example 25.**

1-(4-Methoxybenzoyl-D,L-2-aminothiazol-4-ylglyciny)-4-[(piperidin-1-yl)methyl]piperidine Hydrochloride Salt.

The title compound was prepared by the coupling of

1-(D,L-2-aminothiazol-4-ylglyciny)-4-[(piperidin-1-yl)-methyl]piperidine and 4-methoxybenzoic acid using a procedure similar to that described in Example 24.

^1NMR

LCMS, m/e 472 (M+1)

25 **Example 26.**

1-(4-Methoxybenzoyl-D,L-2-methylthiazol-4-ylglyciny)-4-[(piperidin-1-yl)methyl]piperidine Hydrochloride Salt.

The title compound was prepared by the coupling of

1-(D,L-2-methylthiazol-4-ylglyciny)-4-[(piperidin-1-yl)-methyl]piperidine and 4-methoxybenzoic acid using a procedure similar to that described in Example 24.

^1NMR

LCMS, m/e 471 (M+1)

Example 27.

1-(3-Chloroindole-6-carbonyl-D-phenylglyciny)-4-[(cyclohexylamino)methyl]piperidine Hydrochloride.

- 5 **A. 1-(1-Methylsulfonyl-3-chloroindole-6-carbonyl-D-phenylglyciny)-4-[(cyclohexylamino)methyl]piperidine.**

Using a procedure similar to that described above for Example 6, 1-(1-methylsulfonyl-3-chloroindole-6-carbonyl-D-phenylglyciny)-4-(methylsulfonyloxymethyl)piperidine (940 mg, 10 1.6 mmol) and cyclohexylamine (1.3 mL, 11.2 mmol) afforded, after purification by column chromatography (SiO₂: 2% 2 N ammonia in methanol:methylene chloride), 600 mg (63%) of the title compound.

¹NMR

- 15 IS-MS, m/e 585 (M+1)

B. 1-(3-Chloroindole-6-carbonyl-D-phenylglyciny)-4-[(cyclohexylamino)methyl]piperidine Hydrochloride.

To a solution of 1-(1-methylsulfonyl-3-chloroindole-6-carbonyl-D-phenylglyciny)-4-[(cyclohexylamino)methyl]- 20 piperidine (100 mg, 0.17 mmol) in 4 mL of methanol was added potassium hydroxide (50 mg, 0.89 mmol). After stirring for 1 h, filtration of the resultant white precipitate afforded 35 mg (40%) of the title compound.

25 ¹NMR

IS-MS, m/e 507 (M+1)

The free base was converted into the hydrochloride salt.

Alternatively, using the method described above for 30 Example 6, reaction of Intermediate C-3 and cyclohexylamine affords, after purification and conversion to the hydrochloride salt, the title compound.

Example 28.

1-(Indole-6-carbonyl-D-phenylglyciny1)-4-[(pyrid-4-ylamino)-methyl]piperidine.

To a solution of indole-6-carboxylic acid (144 mg, 0.894 mmol), 1-(D-phenylglyciny1)-4-[(pyrid-4-ylamino)-methyl]piperidine (290 mg, 0.894 mg) and HOAT (134 mg, 0.983 mmol) in DMF (7 mL) was added EDCI (189 mg, 0.983 mmol) and the mixture stirred for 5 h at ambient temperature and 18 h at 5 °C. The DMF was removed in vacuo and the residue
10 purified by RPHPLC. The TFA salt obtained was converted to the free base by dissolving in 2:1 chloroform:isopropanol and washing with aqueous sodium bicarbonate, drying (MgSO₄) and evaporating solvent in vacuo to give the title compound (325 mg, 78%).

15 ¹NMR

The following compounds are prepared using similar procedures to those described above and the requisite starting materials:

20

1-[Indole-6-carbonyl-D,L-(5-thiazoyl)glyciny1]-4-[(piperidin-1-yl)methyl]piperidine

**1-[Indole-6-carbonyl-D,L-(2-aminothiazol-4-yl)glyciny1]-4-
25 [(piperidin-1-yl)methyl]piperidine**

**1-[Indole-6-carbonyl-D,L-(2-methylthiazol-4-yl)glyciny1]-4-
[(piperidin-1-yl)methyl]piperidine**

**30 1-[3-Methylindole-6-carbonyl-D,L-(4-pyridyl)glyciny1]-4-
[(piperidin-1-yl)methyl]piperidine**

**1-[3-Chloroindole-6-carbonyl-D,L-(4-pyridyl)glyciny1]-4-
[(piperidin-1-yl)methyl]piperidine**

1- [3-Methylindole-6-carbonyl-D,L- (2-methoxyphenyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

5 1- [3-Chloroindole-6-carbonyl-D,L- (2-methoxyphenyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [Indole-6-carbonyl-D,L- (1-naphthyl) glyciny] -4- [(piperidin-
1-yl)methyl]piperidine

10 1- [3-Methylindole-6-carbonyl-D,L- (1-naphthyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [3-Chloroindole-6-carbonyl-D,L- (1-naphthyl) glyciny] -4-
15 [(piperidin-1-yl)methyl]piperidine

1- [Indole-6-carbonyl-D,L- (2-chlorophenyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

20 1- [3-Methylindole-6-carbonyl-D,L- (2-chlorophenyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [3-Chloroindole-6-carbonyl-D,L- (2-chlorophenyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

25 1- [Indole-6-carbonyl-D,L- (8-quinolinyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [3-Methylindole-6-carbonyl-D,L- (8-quinolinyl) glyciny] -4-
30 [(piperidin-1-yl)methyl]piperidine

1- [3-Chloroindole-6-carbonyl-D,L- (8-quinolinyl) glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [Indole-6-carbonyl-D,L- (4-quinolinyl)glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

1- [3-Methylindole-6-carbonyl-D,L- (4-quinolinyl)glyciny] -4-
5 [(piperidin-1-yl)methyl]piperidine

1- [3-Chloroindole-6-carbonyl-D,L- (4-quinolinyl)glyciny] -4-
[(piperidin-1-yl)methyl]piperidine

10 Assay protocols

Enzyme Inhibition assays:

The ability of a test compound to inhibit factor Xa may be
15 evaluated in one or more of the following Enzyme Inhibition
assays, or in other standard assays known to those skilled in
the art.

Enzyme Inhibition Assay 1

20

Enzyme assays were carried out at room temperature in 0.1M
phosphate buffer, pH7.4 according to the method of Tapparelli
et al (J. Biol. Chem. 1993,268,4734-4741). Purified human
factor Xa, trypsin, thrombin and plasmin were purchased from
25 Alexis Corporation, Nottingham, UK. Urokinase was purchased
from Calbiochem, Nottingham, UK. Chromogenic substrates for
these enzymes; pefachrome-FXA, pefachrome-TRY, pefachrome-TH,
pefachrome-PL and pefachrome-UK were purchased from Pentapharm
AG, Basel, Switzerland. Product (p-nitroaniline) was
30 quantified by adsorption at 405nm in 96 well microplates using
a Dynatech MR5000 reader (Dynex Ltd, Billingshurst, UK). Km
and Ki were calculated using SAS PROC NLIN (SAS Institute,
Cary, NC, USA, Release 6.11) K_m values were determined as
100.9µM for factor Xa/pefachrome-FXA and 81.6µM for

trypsin/pefachrome-TRY. Inhibitor stock solutions were prepared at 40mM in Me2SO and tested at 500µM, 50µM and 5µM. Accuracy of Ki measurements was confirmed by comparison with Ki values of known inhibitors of factor Xa and trypsin.

5

In agreement with published data, benzamidine inhibited factor Xa, trypsin, thrombin, plasmin and urokinase with Ki values of 155µM, 21µM, 330nM, 200nM and 100nM respectively. NAPAP inhibited thrombin with a Ki value of 3nM. Compounds of the

10 invention were found to have activity in these assays.

Enzyme Inhibition Assay 2

Human factor Xa and human thrombin were purchased from Enzyme
15 Research Laboratories (South Bend, Indiana, USA). Other proteases were from other commercial sources. Chromogenic para-nitroanilide peptide protease substrates were purchased from Midwest Biotech (Fishers, Indiana, USA).

20 The binding affinities for human factor Xa were measured as apparent association constants (K_{ass}) derived from protease inhibition kinetics as described previously.^{a,b,c,d} The apparent K_{ass} values were obtained using automated (BioMek-1000) dilutions of inhibitors (K_{ass} determinations are
25 performed in triplicate at each of four-eight inhibitor concentrations) into 96-well plates and chromogenic substrate hydrolysis rates determined at 405 nm using a Thermomax plate reader from Molecular Devices (San Francisco). For factor Xa inhibition, the assay protocol was: 50 µl buffer (0.06 M tris,
30 0.3 M NaCl, pH 7.4); 25 µl inhibitor test solution (in MeOH); 25 µl human factor Xa (32 nM in 0.03 M tris, 0.15 M NaCl, 1 mg/ml HSA); finally, 150 µl BzIleGluGlyArgpNA (0.3 mM in water) added within 2 min to start hydrolysis. Final factor Xa was

3.2 nM. Free [Xa] and bound [Xa] were determined from linear standard curves on the same plate by use of SoftmaxPro software for each inhibitor concentration and apparent Kass calculated for each inhibitor concentration which produced

5 hydrolysis inhibition between 20% and 80% of the control (3.2 nM factor Xa): apparent Kass = $[E:I]/[E_f][I_f] = [E_b]/[E_f][I^0-I_b]$. The apparent Kass values so obtained are

approximately the inverse of the Ki for the respective inhibitors $[1/appKass = app Ki]$. The variability of mean

10 apparent Kass values determined at the single substrate concentration was +/- 15%. The assay system Km was measured as 0.347 +/- 0.031 mM [n=4]; and Vmax was 13.11 +/- 0.76 μ M/min.

15 Kass values were determined with thrombin and other proteases using the same protocol with the following enzyme and substrate concentrations: thrombin 5.9 nM with 0.2 mM BzPheValArgpNA; XIa 1.2 nM with 0.4 mM pyroGluProArgpNA; XIIa 10 nM with 0.2 mM HDProPheArgpNA; plasmin 3.4 nM with 0.5 mM
20 HDValLeuLyspNA; nt-PA 1.2 nM with 0.8 mM HDileProArgpNA; and urokinase 0.4 nM with 0.4 mM pyroGluGlyArgpNA; aPC 3 nM with 0.174 mM pyroGluProArgpNA; plasma kallikrein 1.9 nM with D-PropPheArgpNA; bovine trypsin 1.4 nM with 0.18 mM BzPheValArgpNA.

25

Citations

(a) Sall DJ, JA Bastian, SL Briggs, JA Buben, NY Chirgadze, DK Clawson, ML Denny, DD Giera, DS Gifford-
30 Moore, RW Harper, KL Hauser, VJ Klimkowski, TJ Kohn, H-S Lin, JR McCowan, AD Palkowitz, GF Smith, ME Richett, K Takeuchi, KJ Thrasher, JM Tinsley, BG Utterback, S-CB Yan, M Zhang. Dibasic Benzo[b]thiophenes Derivatives as a Novel Class of Active Site Directed Thrombin

Inhibitors. 1. Determination of the Serine Protease Selectivity, Structure-Activity Relationships and Binding Orientation. J Med Chem 40 3489-3493 (1997).

- 5 (b) Smith GF, TJ Craft, DS Gifford-Moore, WJ Coffman, KD Kurz, E Roberts, RT Shuman, GE Sandusky, ND Jones, N Chirgadze, and CV Jackson. A Family of Arginal Thrombin Inhibitors Related to Efegatran. Sem. Thrombos. Hemost. 22, 173-183 (1996).
- 10 (c) Smith GF, DS Gifford-Moore, TJ Craft, N Chirgadze, KJ Ruterbories, TD Lindstrom, JH Satterwhite. Efegatran: A New Cardiovascular Anticoagulant. In New Anticoagulants for the Cardiovascular Patient. Ed. R Pifarre. Hanley & Belfus, Inc., Philadelphia (1997) pp 265-300.
- 15 (d) Sall DJ, JA Bastian, NY Chirgadze, ML Denny, MJ Fisher, DS Gifford-Moore, RW Harper, VJ Klimkowski, TJ Kohn, HS Lin, JR McCowan, ME Richett, GF Smith, K Takeuchi, JE Toth, M Zhang. Diamino Benzo[b]thiophene
- 20 Derivatives as a Novel Class of Active Site Directed Thrombin Inhibitors: 5. Potency, Efficacy and Pharmacokinetic Properties of Modified C-3 Side Chain Derivatives. In press, J Med Chem (1999).
- 25 In general, the compounds of formula (I) exemplified herein have been found to exhibit a K_i of 10 μM or less in Assay 1 and/or a K_{ass} of at least 0.1×10^6 L/mole in Assay 2.
- The ability of a test compound to elongate Partial
- 30 Thromboplastin Time (Prothrombin Time) may be evaluated in the following test protocols.

Partial Thromboplastin Time (Prothrombin) Test Protocol

Venous blood was collected into 3.2% (0.109m) trisodium citrate vacutainer tubes at 1 volume of anticoagulant to nine volumes of blood. The blood cells were separated by centrifugation at 700g for ten minutes to yield plasma, which 5 was frozen at 70°C until required.

To perform the test, 100µl of plasma was pipetted into in a glass test tube, 1µl of test compound in DMSO was added, and allowed to warm to 37° over two minutes. 100µl of warm (37°) Manchester (tissue thromboplastin) reagent (Helena Biosciences, 10 UK) was added, allowed to equilibrate for two minutes. 100µl of warm (37°) 25mM calcium chloride solution was added to initiate clotting. The test tube was tilted three times through a 90° angle every five seconds to mix the reagents and the time to clot formation recorded. Data from a series of 15 observations and test compound concentrations are analysed by a SAS statistical analysis program and a CT2 (Concentration required to double clotting time) for each compound is generated.

20 Compounds of the invention were found to significantly elongate the partial thromboplastin time (Prothrombin time).

Alternative Prothrombin Time and APTT Protocols

25 Coagulation Determinations. Prothrombin Times and APTT values were determined in HUMAN PLASMA with a STA instrument (Stago).

BioPT is a special non-plasma clotting assay triggered with human tissue factor (Innovin). Possible binding to albumen or to lipid was assessed by comparing the BioPT effects in the 30 presence/absence of 30 mg/ml human albumen (HSA) and 1 mg/ml phosphatidyl choline (PC). Inhibitors were delivered in 50% MeOH vehicle.

APTT ASSAY

75 µl plasma Citrol *Baxter-Dade* Citrated Normal

Human Plasma

25 µl test sol'n

5 75 µl Actin *Baxter-Dade* Activated Cephaloplastin incubate 2 min
min. @ 37°

75 µl CaCl₂ (0.02 M)

PT ASSAY

10 75 µl plasma

25 µl test sol'n

75 µl saline incubate 1 min. @ 37° C

75 µl Innovin *Baxter-Dade* Recombinant Human Tissue Factor

15 Compounds of the invention were found to be potent inhibitors
of factor Xa.